

Accepted Manuscript

Function of the head–tail junction in the activity of myosin II


Takeshi Haraguchi, Kei Honda, Yuichi Wanikawa, Nao Shoji, Keiichi Yamamoto, Kohji Ito

PII: S0006-291X(13)01509-X

DOI: <http://dx.doi.org/10.1016/j.bbrc.2013.09.038>

Reference: YBBRC 30869

To appear in: *Biochemical and Biophysical Research Communications*



Please cite this article as: T. Haraguchi, K. Honda, Y. Wanikawa, N. Shoji, K. Yamamoto, K. Ito, Function of the head–tail junction in the activity of myosin II, *Biochemical and Biophysical Research Communications* (2013), doi: <http://dx.doi.org/10.1016/j.bbrc.2013.09.038>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Function of the head–tail junction in the activity of myosin II

Takeshi Haraguchi*, Kei Honda*, Yuichi Wanikawa, Nao Shoji, Keiichi Yamamoto,
and Kohji Ito[†]

Department of Biology, Graduate School of Science, Chiba University, Inage-ku, Chiba
263-8522, Japan

[†]Corresponding author at Department of Biology, Graduate School of Science, Chiba
University, Inage-ku, Chiba 263-8522, Japan.

E-mail address: k-ito@faculty.chiba-u.jp

TEL/FAX: +81-43-290-2812

*These authors contributed equally to this work.

Abbreviations used: HMM, heavy meromyosin; S1, subfragment-1; WT, wild type

Download English Version:

<https://daneshyari.com/en/article/8297645>

Download Persian Version:

<https://daneshyari.com/article/8297645>

[Daneshyari.com](https://daneshyari.com)