Complex II phosphorylation is triggered by unbalanced redox homeostasis in cells lacking complex III

Concetta Valentina Tropeano, Jessica Fiori, Valerio Carelli, Leonardo Caporali, Fevzi Daldal, Anna Maria Ghelli, Michela Rugolo

PII: S0005-2728(17)30194-9
DOI: https://doi.org/10.1016/j.bbabio.2017.12.003
Reference: BBABIO 47857

To appear in:

Received date: 4 September 2017
Revised date: 11 December 2017
Accepted date: 16 December 2017

Please cite this article as: Concetta Valentina Tropeano, Jessica Fiori, Valerio Carelli, Leonardo Caporali, Fevzi Daldal, Anna Maria Ghelli, Michela Rugolo , Complex II phosphorylation is triggered by unbalanced redox homeostasis in cells lacking complex III. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Bbabio(2017), https://doi.org/10.1016/j.bbabio.2017.12.003

Complex II phosphorylation is triggered by unbalanced redox homeostasis in cells lacking complex III

Concetta Valentina Tropeano[1], Jessica Fiori[1], Valerio Carelli[2,3], Leonardo Caporali[3], Fevzi Daldal[4], Anna Maria Ghelli[1] and Michela Rugolo[1*]*

[1]Dipartimento di Farmacia e Biotecnologie (FABIT), Università di Bologna, Bologna, Italy;
[2] IRCCS Istituto delle Scienze Neurologiche di Bologna (ISNB), Ospedale Bellaria, Bologna, Italy;
[3]Unità di Neurologia, Dipartimento di Scienze Biomediche e Neuromotorie (DIBINEM), Università di Bologna, Bologna, Italy.
[4] Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA.

*Corresponding authors:
Michela Rugolo, michela.rugolo@unibo.it; and Anna Maria Ghelli, annamaria.ghelli@unibo.it
Dipartimento di Farmacia e Biotecnologie (FABIT), Università degli studi di Bologna
Lab di Biochimica Cellulare, Via Selmi 3, 40126 Bologna, Italy.