


Accepted Manuscript

Mediator, known as a coactivator, can act in transcription initiation in an activator-independent manner in vivo

Hélène Jean-Jacques, Siew Lay Poh, Laurent Kuras



PII: S1874-9399(18)30189-5
DOI: doi:[10.1016/j.bbagr.2018.07.001](https://doi.org/10.1016/j.bbagr.2018.07.001)
Reference: BBAGRM 1267

To appear in: *BBA - Gene Regulatory Mechanisms*

Received date: 3 May 2018
Revised date: 28 June 2018
Accepted date: 4 July 2018

Please cite this article as: Hélène Jean-Jacques, Siew Lay Poh, Laurent Kuras , Mediator, known as a coactivator, can act in transcription initiation in an activator-independent manner in vivo. *Bbagrm* (2018), doi:[10.1016/j.bbagr.2018.07.001](https://doi.org/10.1016/j.bbagr.2018.07.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Mediator, known as a coactivator, can act in transcription initiation in an activator-independent manner *in vivo*

Hélène Jean-Jacques, Siew Lay Poh and Laurent Kuras*

Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ. Paris-Sud, Université Paris-Saclay, Gif-sur-Yvette, France

*Corresponding author: Institute for Integrative Biology of the Cell (I2BC), Avenue de la Terrasse, 91198 Gif-sur-Yvette cedex, France.

E-mail address: laurent.kuras@i2bc.paris-saclay.fr; Tel. +33169823159

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/8300251>

Download Persian Version:

<https://daneshyari.com/article/8300251>

[Daneshyari.com](https://daneshyari.com)