


Accepted Manuscript

Histone variants H3.3 and H2A.Z are incorporated into the β -globin locus during transcription activation via different mechanisms

Jin Kang, Yea Woon Kim, AeRi Kim



PII: S1874-9399(18)30082-8
DOI: doi:[10.1016/j.bbagr.2018.05.005](https://doi.org/10.1016/j.bbagr.2018.05.005)
Reference: BBAGRM 1255
To appear in: *BBA - Gene Regulatory Mechanisms*
Received date: 23 February 2018
Revised date: 21 May 2018
Accepted date: 31 May 2018

Please cite this article as: Jin Kang, Yea Woon Kim, AeRi Kim , Histone variants H3.3 and H2A.Z are incorporated into the β -globin locus during transcription activation via different mechanisms. Bbagrm (2018), doi:[10.1016/j.bbagr.2018.05.005](https://doi.org/10.1016/j.bbagr.2018.05.005)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Histone variants H3.3 and H2A.Z are incorporated into the β -globin locus during transcription activation via different mechanisms

Jin Kang, Yea Woon Kim, AeRi Kim*

*Department of Molecular Biology, College of Natural Sciences,
Pusan National University, Busan 46241, Republic of Korea*

Keywords: H2A.Z, H3.3, transcription, enhancer, promoter

* To whom correspondence should be addressed. Tel: 82-51-510-3683; Fax: 82-51-513-9258;
Email: kimaeri@pusan.ac.kr

Download English Version:

<https://daneshyari.com/en/article/8300267>

Download Persian Version:

<https://daneshyari.com/article/8300267>

[Daneshyari.com](https://daneshyari.com)