Accepted Manuscript



CaMKK2 kinase domain interacts with the autoinhibitory region through the N-terminal lobe including the RP insert

Salome Kylarova, Katarina Psenakova, Petr Herman, Veronika Obsilova, Tomas Obsil

PII: S0304-4165(18)30219-8
DOI: doi:10.1016/j.bbagen.2018.07.025
Reference: BBAGEN 29174

To appear in: *BBA - General Subjects*

Received date: 14 May 2018
Revised date: 18 July 2018
Accepted date: 22 July 2018

Please cite this article as: Salome Kylarova, Katarina Psenakova, Petr Herman, Veronika Obsilova, Tomas Obsil , CaMKK2 kinase domain interacts with the autoinhibitory region through the N-terminal lobe including the RP insert. Bbagen (2018), doi:10.1016/j.bbagen.2018.07.025

CaMKK2 kinase domain interacts with the autoinhibitory region through the N-terminal lobe including the RP insert

Salome Kylarova[1,2], Katarina Psenakova[1,2], Petr Herman[3], Veronika Obsilova[2], Tomas Obsil[1,2]

[1]Department of Physical and Macromolecular Chemistry, Faculty of Science, Charles University, Prague, Czech Republic

[2]BioCeV – Institute of Physiology, The Czech Academy of Sciences, Vestec, Czech Republic

[3]Institute of Physics, Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic

Correspondence

T. Obsil, Department of Physical and Macromolecular Chemistry, Faculty of Science, Charles University, Prague, Czech Republic

Tel: +420 221951303

E-mail: obsil@natur.cuni.cz

V. Obsilova, BioCeV – Institute of Physiology, The Czech Academy of Sciences, Prague, Czech

Republic

Tel: +420 325873513

E-mail: veronika.obsilova@fgu.cas.cz