

Accepted Manuscript


Cystathionine γ lyase–hydrogen sulfide increases peroxisome proliferator-activated receptor γ activity by sulphydration at C139 site thereby promoting glucose uptake and lipid storage in adipocytes

Junyan Cai, Xiaoqin Shi, Huamin Wang, Jinghui Fan, Yongliang Feng, Xianjuan Lin, Jichun Yang, Qinghua Cui, Chaoshu Tang, Guoheng Xu, Bin Geng

PII: S1388-1981(16)30052-X
DOI: doi: [10.1016/j.bbalip.2016.03.001](https://doi.org/10.1016/j.bbalip.2016.03.001)
Reference: BBAMCB 57920

To appear in: *BBA - Molecular and Cell Biology of Lipids*

Received date: 29 December 2015
Revised date: 25 February 2016
Accepted date: 1 March 2016



Please cite this article as: Junyan Cai, Xiaoqin Shi, Huamin Wang, Jinghui Fan, Yongliang Feng, Xianjuan Lin, Jichun Yang, Qinghua Cui, Chaoshu Tang, Guoheng Xu, Bin Geng, Cystathionine γ lyase–hydrogen sulfide increases peroxisome proliferator-activated receptor γ activity by sulphydration at C139 site thereby promoting glucose uptake and lipid storage in adipocytes, *BBA - Molecular and Cell Biology of Lipids* (2016), doi: [10.1016/j.bbalip.2016.03.001](https://doi.org/10.1016/j.bbalip.2016.03.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Cystathionine γ lyase–hydrogen sulfide increases peroxisome
proliferator-activated receptor γ activity by sulfhydration at C139 site
thereby promoting glucose uptake and lipid storage in adipocytes

Junyan Cai^{a*}, Xiaoqin Shi^{b*}, Huamin Wang^a, Jinghui Fan^a, Yongliang Feng^a, Xianjuan Lin^a,
Jichun Yang^{a,c}, Qinghua Cui^{a,c}, Chaoshu Tang^a, Guoheng Xu^a, Bin Geng^{a,c,d}

- a. Department of Physiology and Pathophysiology, Peking University Health Science Center, P.R. China.
- b. Department of Physiology, Hebei University Health Science Center, P.R. China.
- c. Center for Noncoding RNA Medicine, Peking University Health Science Center, P.R. China.
- d. Hypertension Center, Fuwai Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College

* These authors contributed equally to this work

Corresponding Author.

Bin Geng, Department of Physiology and Pathophysiology, Peking University Health Science Center, P.R. China. NO. 38 Xuyuan Road, Haidian District, Bei Jing, China. Tel: +86-010-82802474; Fax: +86-010-82805613; E-mail: bingeng@hsc.pku.edu.cn.

Running title:

CSE–H₂S induces PPAR γ sulfhydration

Conflicts of interest: The authors declare no conflicts of interest.

Manuscript statistics:

Total words:	5875 Words
Reference numbers:	41
Number of figures:	7
Number of tables:	2

Download English Version:

<https://daneshyari.com/en/article/8301724>

Download Persian Version:

<https://daneshyari.com/article/8301724>

[Daneshyari.com](https://daneshyari.com)