

Accepted Manuscript

IP-receptor and PPARs trigger the conversion of human white to brite adipocyte induced by carbaprostacyclin

Rayane A. Ghandour, Maude Giroud, Alexandros Vegiopoulos, Stephan Herzig, Gérard Ailhaud, Ez-Zoubir Amri, Didier F. Pisani

PII: S1388-1981(16)00008-1
DOI: doi: [10.1016/j.bbalip.2016.01.007](https://doi.org/10.1016/j.bbalip.2016.01.007)
Reference: BBAMCB 57880

To appear in: *BBA - Molecular and Cell Biology of Lipids*

Received date: 8 October 2015
Revised date: 4 January 2016
Accepted date: 12 January 2016



Please cite this article as: Rayane A. Ghandour, Maude Giroud, Alexandros Vegiopoulos, Stephan Herzig, Gérard Ailhaud, Ez-Zoubir Amri, Didier F. Pisani, IP-receptor and PPARs trigger the conversion of human white to brite adipocyte induced by carbaprostacyclin, *BBA - Molecular and Cell Biology of Lipids* (2016), doi: [10.1016/j.bbalip.2016.01.007](https://doi.org/10.1016/j.bbalip.2016.01.007)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

IP-receptor and PPARs trigger the conversion of human white to brite adipocyte induced by carbaprostacyclin

Rayane A. Ghandour^{1,2,3}, Maude Giroud^{1,2,3}, Alexandros Vegiopoulos⁴, Stephan Herzig^{5,6,7,8}, Gérard Ailhaud^{1,2,3}, Ez-Zoubir Amri^{1,2,3*} and Didier F. Pisani^{1,2,3*}

¹Univ. Nice Sophia Antipolis, iBV, UMR 7277, 06100 Nice, France.

²CNRS, iBV UMR 7277, 06100 Nice, France.

³Inserm, iBV, U1091, 06100 Nice, France.

⁴DKFZ Junior Group Metabolism and Stem Cell Plasticity, German Cancer Research Center, Heidelberg, Germany,

⁵Institute for Diabetes and Cancer (IDC), Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany

⁶Joint Heidelberg-IDC Translational Diabetes Program, Heidelberg University Hospital, Heidelberg, Germany

⁷Molecular Metabolic Control, Medical Faculty, Technical University Munich, Germany

⁸German Center for Diabetes Research (DZD), Neuherberg, Germany

*Corresponding authors: Dr. Didier Pisani, iBV; Université de Nice Sophia-Antipolis, UMR7277 CNRS - UMR1091 INSERM; Faculté de Médecine; 28 Avenue de Valombrose 06107 Nice cedex 2, France; Tel: +33 493 37 77 07; E-mail: pisani@unice.fr and Dr. Ez-Zoubir Amri, Tel: +33 493 37 70 82; e-mail: amri@unice.fr,

Author contributions. The author(s) have made the following declarations about their contributions: Conceived and designed the experiments: DP, EZA. Performed the experiments: RAG, MG, AV, DFP. Analyzed the data: RAG, AV, SH, GA, DFP, EZA. Wrote the manuscript: RAG, GA, DFP, EZA.

Acknowledgements. This work was supported by CNRS, EU FP7 project DIABAT (HEALTH-F2-2011-278373) and Nutricia Research Foundation ("2011-25"). RAG is a recipient of a fellowship from the Association of Specialization and Scientific Orientation (Lebanon) and Ghazieh township (Lebanon).

Download English Version:

<https://daneshyari.com/en/article/8301755>

Download Persian Version:

<https://daneshyari.com/article/8301755>

[Daneshyari.com](https://daneshyari.com)