


## Accepted Manuscript

Contribution of the exosome-associated form of secreted endoplasmic reticulum aminopeptidase 1 to exosome-mediated macrophage activation

Yoshikuni Goto, Yuko Ogawa, Hiroki Tsumoto, Yuri Miura, Takahiro J. Nakamura, Kenji Ogawa, Yoshihiro Akimoto, Hayato Kawakami, Tamao Endo, Ryohei Yanoshita, Masafumi Tsujimoto



PII: S0167-4889(18)30044-2  
DOI: doi:[10.1016/j.bbamcr.2018.03.009](https://doi.org/10.1016/j.bbamcr.2018.03.009)  
Reference: BBAMCR 18258

To appear in:

Received date: 12 December 2017  
Revised date: 26 February 2018  
Accepted date: 15 March 2018

Please cite this article as: Yoshikuni Goto, Yuko Ogawa, Hiroki Tsumoto, Yuri Miura, Takahiro J. Nakamura, Kenji Ogawa, Yoshihiro Akimoto, Hayato Kawakami, Tamao Endo, Ryohei Yanoshita, Masafumi Tsujimoto, Contribution of the exosome-associated form of secreted endoplasmic reticulum aminopeptidase 1 to exosome-mediated macrophage activation. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Bbamcr*(2018), doi:[10.1016/j.bbamcr.2018.03.009](https://doi.org/10.1016/j.bbamcr.2018.03.009)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Contribution of the exosome-associated form of secreted endoplasmic reticulum  
aminopeptidase 1 to exosome-mediated macrophage activation**

Yoshikuni Goto<sup>a\*</sup>, Yuko Ogawa<sup>a</sup>, Hiroki Tsumoto<sup>b</sup>, Yuri Miura<sup>b</sup>, Takahiro J Nakamura<sup>c</sup>,  
Kenji Ogawa<sup>d</sup>, Yoshihiro Akimoto<sup>e</sup>, Hayato Kawakami<sup>e</sup>, Tamao Endo<sup>b</sup>, Ryohei  
Yanoshita<sup>a</sup>, and Masafumi Tsujimoto<sup>a</sup>

<sup>a</sup> Faculty of Pharmaceutical Sciences, Teikyo-Heisei University, Nakano, Tokyo, 164-8530, Japan, <sup>b</sup> Research Team for Mechanism of Aging, Tokyo Metropolitan Institute of Gerontology, Itabashi, Tokyo 173-0015, Japan., <sup>c</sup> Department of Life Sciences, School of Agriculture, Meiji University, Kawasaki, Kanagawa 214-8571, Japan, <sup>d</sup> Chemical Genetics Laboratory, RIKEN, Wako, Saitama, 351-0198, Japan, <sup>e</sup> Department of Anatomy, Kyorin University School of Medicine, Mitaka, Tokyo 181-8611, Japan

\* Corresponding author at: Faculty of Pharmaceutical Sciences, Teikyo-Heisei University, 4-21-2 Nakano, Nakano, Tokyo, 164-8530, Japan, Telephone: (+81) 3-5860-4259; E-mail: y.goto@thu.ac.jp (Y. Goto)

**Keywords:** aminopeptidase, exosome, macrophages, phagocytosis, nitric oxide, cytokines

Download English Version:

<https://daneshyari.com/en/article/8303665>

Download Persian Version:

<https://daneshyari.com/article/8303665>

[Daneshyari.com](https://daneshyari.com)