

Accepted Manuscript

Title: The MLH1 ATPase Domain is needed for Suppressing Aberrant Formation of Interstitial Telomeric Sequences

Authors: Pingping Jia, Weihang Chai

PII: S1568-7864(18)30025-9
DOI: <https://doi.org/10.1016/j.dnarep.2018.03.002>
Reference: DNAREP 2476

To appear in: *DNA Repair*

Received date: 9-2-2018
Revised date: 2-3-2018
Accepted date: 5-3-2018



Please cite this article as: Pingping Jia, Weihang Chai, The MLH1 ATPase Domain is needed for Suppressing Aberrant Formation of Interstitial Telomeric Sequences, DNA Repair <https://doi.org/10.1016/j.dnarep.2018.03.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Ms: DNAREP_2018_16 R1

**The MLH1 ATPase Domain is needed for Suppressing Aberrant Formation of Interstitial
Telomeric Sequences**

Pingping Jia, Weihang Chai *

Department of Biomedical Sciences, Elson S. Floyd College of Medicine, Washington State
University

Running title: MLH1 ATPase suppresses telomeric sequences insertion

Key words: MLH1, ATPase, interstitial telomere sequences

*Corresponding author:

Weihang Chai, Ph.D., PO Box 1495, Spokane, WA 99210, Tel: 509-358-7575, Fax: 509-358-
7882. Email: wchai@wsu.edu

Download English Version:

<https://daneshyari.com/en/article/8320362>

Download Persian Version:

<https://daneshyari.com/article/8320362>

[Daneshyari.com](https://daneshyari.com)