Anti-inflammatory effect of low molecular weight fucoidan from Saccharina japonica on atherosclerosis in apoE-knockout mice

Yingjie Xu, Jie Xu, Keli Ge, Qingwu Tian, Peng Zhao, Yunliang Guo

PII: S0141-8130(18)31126-7
DOI: doi:10.1016/j.ijbiomac.2018.06.054
Reference: BIOMAC 9894

To appear in: *International Journal of Biological Macromolecules*

Received date: 9 March 2018
Revised date: 5 June 2018
Accepted date: 11 June 2018

Please cite this article as: Yingjie Xu, Jie Xu, Keli Ge, Qingwu Tian, Peng Zhao, Yunliang Guo , Anti-inflammatory effect of low molecular weight fucoidan from Saccharina japonica on atherosclerosis in apoE-knockout mice. Biomac (2017), doi:10.1016/j.ijbiomac.2018.06.054

Anti-inflammatory effect of low molecular weight fucoidan from *Saccharina japonica* on atherosclerosis in apoE-knockout mice

Yingjie Xu[a], Jie Xu[a], Keli Ge [a], Qingwu Tian[b], Peng Zhao[b], Yunliang Guo[a*]

a Institute of Cerebrovascular Diseases, Affiliated Hospital of Qingdao University, Qingdao 266003, China

b Clinical laboratory, Affiliated Hospital of Qingdao University, Qingdao 266003, China

*Corresponding author:

Yunliang Guo, guoqdsd@163.com; Tel/fax: +86-532-8299-1711.