Accepted Manuscript

Food restriction followed by refeeding with a casein- or whey-based diet differentially affects the gut microbiota of pre-pubertal male rats

Majdi Masarwi, Hadas Isaac Solnik, Moshe Phillip, Sima Yaron, Raanan Shamir. Metsada Pasmanic-Chor, Galia Gat-Yablonski

PII: S0955-2863(17)30147-X

DOI: doi: 10.1016/j.jnutbio.2017.08.014

Reference: JNB 7832

To appear in: The Journal of Nutritional Biochemistry

Received date: 13 February 2017 Revised date: 20 August 2017 Accepted date: 29 August 2017



Please cite this article as: Masarwi Majdi, Solnik Hadas Isaac, Phillip Moshe, Yaron Sima, Shamir Raanan, Pasmanic-Chor Metsada, Gat-Yablonski Galia, Food restriction followed by refeeding with a casein- or whey-based diet differentially affects the gut microbiota of pre-pubertal male rats, *The Journal of Nutritional Biochemistry* (2017), doi: 10.1016/j.jnutbio.2017.08.014

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Food restriction followed by refeeding with a casein- or whey-based diet

differentially affects the gut microbiota of pre-pubertal male rats

Majdi Masarwi^{a,b}, Hadas Isaac Solnik^c, Moshe Phillip^{a,b,d}, Sima Yaron^c,

Raanan Shamir^{a,e}, Metsada Pasmanic-Chor^f, Galia Gat-Yablonski^{a,b,d}

^aSackler School of Medicine, Tel Aviv University, Tel Aviv, Israel; ^bFelsenstein Medical Research

Center, Petach Tikva, Israel; ^cFaculty of Biotechnology and Food Engineering, Technion -- Israel

Institute of Technology, Haifa, Israel; ^dThe Jesse Z and Sara Lea Shafer Institute for Endocrinology

and Diabetes, National Center for Childhood Diabetes, Schneider Children's Medical Center of

Israel, Petach Tikva, Israel; eInstitute for Gastroenterology, Nutrition and Liver Diseases, Schneider

Children's Medical Center of Israel, Petach Tikva, Israel; ^fBioinformatics unit, The George S. Wise

Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel

E-mail addresses

M. Masarwi: majdimas@post.tau.ac.il

H. Isaac Solnik: hadasi@efco.co.il

M. Phillip: mosheph@post.tau.ac.il

S. Yaron: simay@tx.technion.ac.il

R. Shamir: shamirraanan@gmail.com

M. Pasmanic-Chor: metsada@post.tau.ac.il

*Corresponding author: Galia Gat-Yablonski, PhD, Institute for Endocrinology and Diabetes,

National Center for Childhood Diabetes, Schneider Children's Medical Center of Israel, Petach

Tikva 4920235, Israel

Tel: +972-3-9376133; Fax: +972-3-9211478; Email: galiagy@post.tau.ac.il

Running title: Diet composition affects gut microbiota

Funding sources: This work was partially supported by the Danone Strauss Research Institute and

the Israeli Dairy Board [grant number 9114].

Conflicts of interest: None.

1

Download English Version:

https://daneshyari.com/en/article/8336454

Download Persian Version:

https://daneshyari.com/article/8336454

<u>Daneshyari.com</u>