



# Corepressors (NCoR and SMRT) as well as coactivators are recruited to positively regulated $1\alpha,25$ -dihydroxyvitamin $D_3$ -responsive genes

Mark B. Meyer, J. Wesley Pike\*

Department of Biochemistry, University of Wisconsin-Madison, Madison, WI 53706, United States

## ARTICLE INFO

### Article history:

Received 12 July 2012

Received in revised form 2 August 2012

Accepted 9 August 2012

### Keywords:

VDR  
ChIP-seq  
Corepressor  
Coactivator  
Coregulator  
Vitamin D

## ABSTRACT

Transcription factors require coactivators and corepressors to modulate transcription in mammalian cells. The vitamin D receptor (VDR) utilizes coactivators and corepressors to gain tight control over the activity of a diverse set of genes that can regulate calcium transport, slow proliferation and promote immune responses. We have recently established the VDR/RXR cisome in human colon cancer cells and have linked these binding sites to the genes that are regulated by  $1\alpha,25$ -dihydroxyvitamin  $D_3$  ( $1,25(OH)_2D_3$ ). In additional studies described herein, we demonstrate that the coactivators SRC1, CBP and MED1 are recruited to upregulated genes to facilitate transcription as expected. SRC1 was the most highly correlated to VDR/RXR binding (50%). However, we also found that corepressor molecules such as NCoR and SMRT were present along with SRC1, CBP or MED1 at these  $1,25(OH)_2D_3$  activated gene enhancers. Interestingly, genome-wide NCoR binding mimicked VDR binding by increasing its association with VDR binding in response to  $1,25(OH)_2D_3$  treatment. Overall, these data indicate a complex role for corepressor and coactivator complexes in the activation or active repression of  $1,25(OH)_2D_3$  responsive genes.

This article is part of a Special Issue entitled 'Vitamin D Workshop'.

© 2012 Elsevier Ltd. All rights reserved.

## 1. Introduction

The steroid hormone  $1\alpha,25$ -dihydroxyvitamin  $D_3$  ( $1,25(OH)_2D_3$ ) tightly regulates calcium and phosphorus homeostasis [1].  $1,25(OH)_2D_3$  acts to regulate genes through its receptor, the vitamin D receptor (VDR), and its heterodimeric partner the retinoid X receptor (RXR). These genes are responsible for retention of calcium in the kidney, absorption of dietary calcium in the intestine and direct remodeling processes across the bone through osteoblasts and bone-resorbing osteoclasts. For the regulation of these processes,  $1,25(OH)_2D_3$  induces genes such as *TRPV6*, *PMCA1B*, and *S100G* whose products are directly involved in the transepithelial transfer of calcium from the gut lumen and *SLC34A1* and *SLC34A3* that facilitate phosphate uptake [2,3] and genes such as *CYP3A4*, *CYP3A7*, *CYP2B6*, and *ABCB1* that function to detoxify lithocholic acid and other secondary bile acids or are involved in the transport and metabolism of foreign compounds [4–6]. It is also known that  $1,25(OH)_2D_3$  exerts activity and regulation on the VDR gene itself as well as its catabolic enzyme *CYP24A1* [7,8].

Coactivators are essential for transcription initiation and are classically believed to provide linkage from receptor complexes to

the basal transcriptional machinery [9]. Indeed it is now known that coactivators have catalytic domains that are responsible for modification of the chromatin environment such as acetylation, methylation, phosphorylation and many others [10]. The modulation of these chromatin marks defines the epigenome that drives cell-type and tissue-type specificity. Currently, there are over 300 transcriptional coregulators that have been described in the literature [11]. Coactivators, those believed to facilitate transcription, and corepressors, those believed to inhibit transcription, have been interchangeably described in the activation and/or repression of genes, making characterization difficult. It has been demonstrated, for example, that for full activation of estrogen responsive genes, the corepressor SMRT is required in the activation complex [12]. The coactivators and corepressors are able to directly interact with nuclear receptors like the VDR through LXXLL protein motifs [13] and SMRT is directly involved with vitamin D-mediated transcription [14] as well as other coregulatory molecules [15]. It is believed that ligand activated receptors can change conformations and preference for coactivators based upon which ligands are receptor bound [16]. Furthermore, post translational modifications of coactivators further diversify the activities transferred during coactivation of genes.

Recent advances in transcription research have revealed an extensive array of instructional epigenetic marks that are inserted across the genome in a cell-type specific manner [17]. Some epigenetic marks are associated specifically with regulatory regions, indicating that cellular phenotype is a direct consequence of the

\* Corresponding author. Department of Biochemistry, University of Wisconsin-Madison, 433 Babcock Drive, Madison, WI 53706, United States. Tel.: +1 608 262 8229; fax: +1 608 263 9609.

E-mail address: [pike@biochem.wisc.edu](mailto:pike@biochem.wisc.edu) (J.W. Pike).

establishment of cell-specific enhancers by early lineage-specific transcription factors [18,19]. Functional binding sites for the VDR/RXR revealed transcription factor interactions as well as the coregulator recruitment processes which identified at least some of the consequences of these interactions at sites on the genome [20]. In the current studies, we expanded on our recent discovery of the VDR/RXR cistrome by analyzing the coactivators (SRC1, CBP, MED1) as well as corepressors (NCoR, SMRT) involved during 1,25(OH)<sub>2</sub>D<sub>3</sub>-mediated transcription. We found a high correlation between SRC1 and CBP occupancy and transcriptional activation; however, we also found that repressors such as NCoR and SMRT were present in the activation complexes. These studies highlight the complex nature of transcription and coregulatory molecules.

## 2. Materials and methods

### 2.1. Reagents

1,25(OH)<sub>2</sub>D<sub>3</sub> was obtained from Tetrionics, Inc. (Madison, WI). Antibodies to SRC-1 (M-341, sc-8495), CBP (A-22, sc-3996), MED1 (M-255, sc-8998), NCoR (C-20, sc-1609) were purchased from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA). SMRT (PA1-843) antibody was purchased from Affinity Bioreagents (Thermo Fisher, Rockford, IL).

### 2.2. Cell culture

Human LS180 CRC cells were obtained from ATCC (Manassas, VA). LS180 cells were cultured in minimum Eagle's medium supplemented with 10% non-heat-inactivated fetal bovine serum from Hyclone (Logan, UT), 1% non-essential amino acids, 1% sodium pyruvate, and 1% penicillin-streptomycin from Invitrogen as previously reported [21].

### 2.3. Chromatin immunoprecipitation sequencing (ChIP-seq)

LS180 cells were treated for 3 h with vehicle or 10<sup>−7</sup> M 1,25(OH)<sub>2</sub>D<sub>3</sub> prior to Chromatin ImmunoPrecipitation which was performed as described previously [21,22]. ChIP-DNA was prepared and amplified using the Illumina ChIP-seq DNA preparation kit (1003473, #11257047 RevA), clusters formed and sequenced on the Illumina GAIIx or HiSeq2000 sequencers by the University of Wisconsin - Madison DNA Sequencing Facility in the University of Wisconsin- Madison Biotechnology Center [22]. Samples were further processed by two methods QuEST and HOMER. Peaks were accepted if they passed criteria of both methods. The data were analyzed with QuEST 2.4 [23] and HOMER [18] as previously reported [22]. All data and tracks have been deposited in the Gene Expression Omnibus (GEO) at GSE39277.

## 3. Results and discussion

### 3.1. VDR/RXR interact with coactivators to modulate transcription

We have previously quantitated the number of DNA binding sites for the vitamin D receptor (VDR) and its heterodimer partner retinoid X receptor (RXR) across the LS180 (human colon adenocarcinoma) genome using ChIP-seq analysis (FDR < 0.001) [22]. Cells were treated with either ethanol vehicle or 1,25(OH)<sub>2</sub>D<sub>3</sub> for 3 h and then subjected to the above analyses using validated antibodies to either VDR or RXR. We determined the number of high confidence peaks for VDR and RXR overlap in the 1,25(OH)<sub>2</sub>D<sub>3</sub> condition to be 638 sites. A mapping analysis of peaks to surrounding genes revealed that 98% of VDR/RXR binding sites were located within

either intergenic or intronic regions, which is consistent with findings for most transcription factors [24–26].

Transcription factor nuclear receptors, such as VDR and RXR, are known to recruit a variety of coregulatory complexes that are essential for altering gene expression [9]. To assess the ability of VDR to recruit these coregulatory complexes to target genes in a genome-wide manner, we conducted individual ChIP-seq analyses in LS180 cells using antibodies to the coactivators SRC1, MED1 and CBP, and to the corepressors NCoR and SMRT following treatment with either vehicle or 1,25(OH)<sub>2</sub>D<sub>3</sub>. Peaks were identified and quantitated as described in [22] using QuEST and HOMER [18,23]. As can be seen in Fig. 1a, while a number of sites for each of the coregulators was present under basal conditions, as might be anticipated for these general coregulators, an overlapping yet unique set of sites for each of these transcription factors was induced by 1,25(OH)<sub>2</sub>D<sub>3</sub> as well. With the exception of those for SMRT, the total number of binding sites for all coregulators increased in the presence of 1,25(OH)<sub>2</sub>D<sub>3</sub>. The number of binding sites for VDR is greatly increased by 1,25(OH)<sub>2</sub>D<sub>3</sub> treatment [22]. These data suggest that 1,25(OH)<sub>2</sub>D<sub>3</sub> likely promotes the recruitment of not only coactivators but also corepressors such as NCoR or SMRT to gene targets as well. We next correlated these coregulatory sites to the peak centers of the 638 1,25(OH)<sub>2</sub>D<sub>3</sub>-induced VDR/RXR-bound sites. In this analysis, SRC1 peak frequency was most highly correlated with VDR/RXR peak frequency (50%) and the peaks were located very near VDR/RXR peak centers. Moreover, we also found that SRC1 tracked most closely with regulated genes of interest suggesting a strong functional relationship between SRC1 and VDR. Only the peak frequency of NCoR binding was reduced in the presence of 1,25(OH)<sub>2</sub>D<sub>3</sub>, suggesting that at some sites ligand activation of the VDR could result in corepressor displacement. Clearly, while corepressors may function at genes to mediate repression by 1,25(OH)<sub>2</sub>D<sub>3</sub>, they may also exert coactivator function as has been observed for the estrogen receptor (ER) [12]. These data suggest that coregulators may interact uniquely at VDR/RXR-bound sites and that these sites may display unique functional activities as well.

Since all 638 VDR/RXR binding sites were annotated to specific genomic locations and to their nearest gene neighbors, we linked these gene loci to the gene expression data documented in our previous work [22]. We then analyzed individual ChIP-seq tag densities for SRC1, CBP, MED1, NCoR and SMRT (25 bp bins) across a 6 kb interval with the VDR/RXR site at its center (±3 kb). As shown in Fig. 1c, each row of tag density represents a different VDR/RXR peak (each peak correlates to nearest gene). Based upon tag density of the coregulators, the data were clustered using hierarchical clustering with centroid linkage and 4 clusters were identified. Data clusters 1 and 4 contained reduced tag densities for many of the coregulators. Indeed, only a few of the genes located adjacent to these VDR/RXR peaks were regulated by 1,25(OH)<sub>2</sub>D<sub>3</sub>. Cluster 2, on the other hand, contained very high tag densities that were associated with all the coregulators and centered specifically on the VDR/RXR peaks themselves. This cluster of sites was linked most closely to the genes highly regulated by 1,25(OH)<sub>2</sub>D<sub>3</sub> such as those listed in Fig. 1d. Cluster 3 represents a small number (eight) of anomalous VDR/RXR binding sites that appeared to be unrelated to gene loci. These data suggest that 1,25(OH)<sub>2</sub>D<sub>3</sub> induces both VDR/RXR binding and coregulator recruitment at specific sites and that these sites are capable of regulating genes that are located nearby.

### 3.2. c-FOS is controlled by coordinated coregulator action

The results of the above studies confirm and extend mechanistic studies conducted over the past decade on several genes known to be regulated by 1,25(OH)<sub>2</sub>D<sub>3</sub> in the intestine or colon, including

Download English Version:

<https://daneshyari.com/en/article/8339258>

Download Persian Version:

<https://daneshyari.com/article/8339258>

[Daneshyari.com](https://daneshyari.com)