

Reproducibility and agreement of different non-invasive methods of endothelial function assessment

Khatera Ibrahimi, Young De Graaf, Richard Draijer, A.H. Jan Danser, Antoinette Maassen VanDenBrink, Anton H. van den Meiracker

PII: S0026-2862(17)30275-3
DOI: https://doi.org/10.1016/j.mvr.2018.01.004
Reference: YMVRE 3755

To appear in: *Microvascular Research*

Received date: 13 December 2017
Revised date: 9 January 2018
Accepted date: 9 January 2018

Please cite this article as: Khatera Ibrahimi, Young De Graaf, Richard Draijer, A.H. Jan Danser, Antoinette Maassen VanDenBrink, Anton H. van den Meiracker , Reproducibility and agreement of different non-invasive methods of endothelial function assessment. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Ymvre(2017), https://doi.org/10.1016/j.mvr.2018.01.004

Reproducibility and agreement of different non-invasive methods of endothelial function assessment

Khatera Ibrahimi[1], Young De Graaf[2], Richard Draijer[2], A.H. Jan Danser[1], Antoinette MaassenVanDenBrink[1], Anton H. van den Meiracker[1]

[1]Division of Pharmacology and Vascular Medicine, Dept. of Internal Medicine, Erasmus MC, Rotterdam, The Netherlands

[2]Unilever R & D, Vlaardingen, The Netherlands

Running title: Thermal Hyperemia and Flow Mediated Dilatation

Sources of funding: This study has been supported by a grant from Unilever R&D, Vlaardingen, The Netherlands

Conflict of Interest: None

Key-words: Flow-mediated dilatation, Local thermal hyperemia, Post-occlusion reactive hyperemia, Nitric oxide.

Word count abstract: 249

Word count text: 2620

Number of references: 27

Number of tables: 4

Number of figures: 1

Correspondence:

Anton H. van den Meiracker,

Division of Pharmacology and Vascular Diseases,

Department of Internal Medicine, Room Ee1402b,

Erasmus MC,

's Gravendeelweg 230,

3015 CE, Rotterdam,

The Netherlands.