Accepted Manuscript



The effect of the electrophilic fatty acid nitro-oleic acid on TRP channel function in sensory neurons

Jonathan M. Beckel, William C. de Groat

PII: S1089-8603(17)30289-6

DOI: 10.1016/j.niox.2018.03.015

Reference: YNIOX 1768

To appear in: *Nitric Oxide*

Received Date: 30 October 2017

Revised Date: 19 March 2018

Accepted Date: 21 March 2018

Please cite this article as: J.M. Beckel, W.C. de Groat, The effect of the electrophilic fatty acid nitro-oleic acid on TRP channel function in sensory neurons, *Nitric Oxide* (2018), doi: 10.1016/j.niox.2018.03.015.

The effect of the electrophilic fatty acid nitro-oleic acid on TRP channel function in sensory neurons

Jonathan M. Beckel[1] and William C. de Groat[1]

[1]Department of Pharmacology and Chemical Biology, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261 USA

Corresponding author:
Jonathan M. Beckel, Ph.D.
W1353 Starzl Biomedical Science Tower
200 Lothrop Street
Pittsburgh, PA 15261
Email: jmbeckel@pitt.edu
Phone: +1-412-383-5004

Keywords: Transient Receptor Potential Vanilloid 1 (TRPV1), Transient Receptor Potential Ankyrin 1 (TRPA1), Transient Receptor Potential Canonical (TRPC), Nitro-oleic acid, afferent nerves, Electrophilic fatty acids, Dorsal root ganglia (DRG), Capsaicin, Allyl isothiocyanate (AITC), Urinary bladder