## Accepted Manuscript

Polysulfides and products of H<sub>2</sub>S/S-nitrosoglutathione in comparison to H<sub>2</sub>S, glutathione and antioxidant Trolox are potent scavengers of superoxide anion radical and produce hydroxyl radical by decomposition of H<sub>2</sub>O<sub>2</sub>

Anton Misak, Marian Grman, Zuzana Bacova, Ingeborg Rezuchova, Sona Hudecova, Elena Ondriasova, Olga Krizanova, Vlasta Brezova, Miroslav Chovanec, Karol Ondrias

PII: S1089-8603(17)30139-8

DOI: 10.1016/j.niox.2017.09.006

Reference: YNIOX 1700

To appear in: Nitric Oxide


Received Date: 22 May 2017

Revised Date: 20 September 2017

Accepted Date: 20 September 2017

Please cite this article as: A. Misak, M. Grman, Z. Bacova, I. Rezuchova, S. Hudecova, E. Ondriasova, O. Krizanova, V. Brezova, M. Chovanec, K. Ondrias, Polysulfides and products of H<sub>2</sub>S/Snitrosoglutathione in comparison to H<sub>2</sub>S, glutathione and antioxidant Trolox are potent scavengers of superoxide anion radical and produce hydroxyl radical by decomposition of H<sub>2</sub>O<sub>2</sub>, *Nitric Oxide* (2017), doi: 10.1016/j.niox.2017.09.006.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



## ACCEPTED MANUSCRIPT

Polysulfides and products of H<sub>2</sub>S/S-nitrosoglutathione in comparison to H<sub>2</sub>S, glutathione and antioxidant Trolox are potent scavengers of superoxide anion radical and produce hydroxyl radical by decomposition of H<sub>2</sub>O<sub>2</sub>

Anton Misak<sup>a,1</sup>, Marian Grman<sup>a,1</sup>, Zuzana Bacova<sup>b</sup>, Ingeborg Rezuchova<sup>c</sup>, Sona Hudecova<sup>a</sup>, Elena Ondriasova<sup>d</sup>, Olga Krizanova<sup>a</sup>, Vlasta Brezova<sup>e</sup>, Miroslav Chovanec<sup>f,2</sup>, Karol Ondrias<sup>a,2,3</sup>

<sup>a</sup> Institute of Clinical and Translational Research, Biomedical Research Center, Slovak Academy of Sciences, Bratislava, Slovak Republic; <sup>b</sup> Institute of Experimental Endocrinology, Biomedical Research Center, Slovak Academy of Sciences, Bratislava, Slovak Republic; <sup>c</sup> Institute of Virology, Biomedical Research Center, Slovak Academy of Sciences, Bratislava, Slovak Republic; <sup>d</sup> Department of Pharmacology and Toxicology, Faculty of Pharmacy, Comenius University, Bratislava, Slovak Republic; <sup>e</sup> Faculty of Chemical and Food Technology, Slovak University of Technology, Bratislava, Slovak Republic; <sup>f</sup> Cancer Research Institute, Biomedical Research Center, Slovak Academy of Sciences, Bratislava, Slovak Republic

Footnotes:

<sup>1</sup>A.M. and M.G. contributed equally to this work.
<sup>2</sup>M.C. and K.O. contributed equally to this work.
<sup>3</sup>To whom correspondence should be addressed. Email: karol.ondrias@savba.sk

Karol Ondrias, Institute of Clinical and Translational Research Biomedical Research Center Slovak Academy of Sciences Dubravska cesta 9 845 05 Bratislava 4 Slovak Republic. Tel.: +421-908577943; e-mail: <u>karol.ondrias@savba.sk</u> Download English Version:

## https://daneshyari.com/en/article/8344595

Download Persian Version:

https://daneshyari.com/article/8344595

Daneshyari.com