Accepted Manuscript

A biochar application protects rice pollen from high-temperature stress

Shah Fahad, Saddam Hussain, Shah Saud, Mohsin Tanveer, Ali Ahsan Bajwa, Shah Hassan, Adnan Noor Shah, Abid Ullah, Chao Wu, Faheem Ahmed Khan, Farooq Shah, Sami Ullah, Yajun Chen, Jianliang Huang

PII: S0981-9428(15)30089-9

DOI: 10.1016/j.plaphy.2015.08.009

Reference: PLAPHY 4261

To appear in: *Plant Physiology and Biochemistry*

Received Date: 8 March 2015

Revised Date: 10 July 2015

Accepted Date: 12 August 2015

Please cite this article as: S. Fahad, S. Hussain, S. Saud, M. Tanveer, A.A. Bajwa, S. Hassan, A.N. Shah, A. Ullah, C. Wu, F.A. Khan, Farooq Shah, Sami Ullah, Y. Chen, J. Huang, A biochar application protects rice pollen from high-temperature stress, Plant Physiology et Biochemistry (2015), doi: 10.1016/j.plaphy.2015.08.009.

A biochar application protects rice pollen from high-temperature stress

Shah Fahad[1], Saddam Hussain[1,2], Shah Saud[3], Mohsin Tanveer[1], Ali Ahsan Bajwa[4], Shah Hassan[5], Adnan Noor Shah[1], Abid Ullah[6], Chao Wu[1], Faheem Ahmed Khan[7], Farooq Shah[1,8], Sami Ullah[9], Yajun Chen[10] and Jianliang Huang[1,11*]

[1]National Key Laboratory of Crop Genetic Improvement, MOA Key Laboratory of Crop Ecophysiology and Farming System in the Middle Reaches of the Yangtze River, College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, Hubei 430070, China

[2]College of Resources and Environment, Huazhong Agricultural University, Wuhan, Hubei 430070, China

[3]Department of Horticulture, Northeast Agricultural University, Harbin 150030, China

[4]Department of Agronomy, University of Agriculture, Faisalabad, Pakistan

[5]Department of Agricultural Extension, Agricultural University Peshawar, Khyber Pakhtunkhwa, Pakistan

[6]National Key Laboratory of Crop Genetic Improvement, Huazhong Agricultural University, Wuhan, P.R. China, 430070

[7]Key Laboratory of Agricultural Animal Genetics, Breeding and Reproduction, Huazhong Agricultural University, Wuhan, 430070, China

[8]Department of Agriculture, Abdul Wali Khan University, Mardan, Khyber Pakhtunkhwa, Pakistan

[9] Department of Botany, Bacha Khan University, Charsadda Khyber Pakhtunkhwa, Pakistan

[10]Horticulture College of Northeast Agricultural University, Harbin, Heilongjiang 150030, China

[11]Hubei Collaborative Innovation Center for Grain Industry, Yangtze University, Hubei, China

***Corresponding author:** Tel. +86 13545874386; Fax: +86 13100633046

E-mail address: jhuang@mail.hzau.edu.cn