Accepted Manuscript

Title: Activation of the salicylic acid signalling pathway in wheat had no significant short-term impact on the diversity of root-associated microbiomes

Authors: Hongwei Liu, Lilia C. Carvalhais, Peer M. Schenk, Paul G. Dennis

PII: S0031-4056(18)30091-X
DOI: https://doi.org/10.1016/j.pedobi.2018.08.001
Reference: PEDOBI 50541

To appear in:

Received date: 9-5-2018
Revised date: 2-8-2018
Accepted date: 6-8-2018

Please cite this article as: Liu H, Carvalhais LC, Schenk PM, Dennis PG, Activation of the salicylic acid signalling pathway in wheat had no significant short-term impact on the diversity of root-associated microbiomes, *Pedobiologia - Journal of Soil Ecology* (2018), https://doi.org/10.1016/j.pedobi.2018.08.001

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Short communication:

Activation of the salicylic acid signalling pathway in wheat had no significant short-term impact on the diversity of root-associated microbiomes

Hongwei Liu[1], Lilia C. Carvalhais[2], Peer M. Schenk[1], Paul G. Dennis[3*]

[1]*School of Agriculture and Food Sciences, The University of Queensland, Brisbane, Queensland 4072, Australia*;

[2]*Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, Ecosciences Precinct, GPO Box 267, Queensland 4001, Australia;*

[3]*School of Earth and Environmental Sciences, The University of Queensland, Brisbane, Queensland 4072, Australia*

*Correspondence: p.dennis@uq.edu.au

Highlights

- The salicylic acid (SA) pathway was activated in wheat grown in two soils.
- Root microbiomes were characterised for control and SA treated plants.
- SA treatment did not significantly affect the diversity of root microbiomes within 72 h.
- SA treatment is unlikely to help engineer beneficial wheat root microbiomes.

Download English Version:

https://daneshyari.com/en/article/8392327

Download Persian Version:

https://daneshyari.com/article/8392327

Daneshyari.com