Accepted Manuscript

Title: Biogeographical patterns of bacterial and archaeal communities from distant hypersaline environments

Authors: M.R. Mora-Ruiz del, A. Cifuentes, F. Font-Verdera, C. Pérez-Fernández, M.E. Farias, B. González, A. Orfila, R. Rosselló-Móra

PII: S0723-2020(17)30178-9
DOI: https://doi.org/10.1016/j.syapm.2017.10.006
Reference: SYAPM 25883

To appear in:

Received date: 10-7-2017
Revised date: 22-10-2017
Accepted date: 23-10-2017

Please cite this article as: M.R.Mora-Ruiz del, A.Cifuentes, F.Font-Verdera, C.Pérez-Fernández, M.E.Farias, B.González, A.Orfila, R.Rosselló-Móra, Biogeographical patterns of bacterial and archaeal communities from distant hypersaline environments, Systematic and Applied Microbiology https://doi.org/10.1016/j.syapm.2017.10.006

Title

Biogeographical patterns of bacterial and archaeal communities from distant hypersaline environments

Authors:

Mora-Ruiz, M. del R.[a], Cifuentes, A.[a], Font-Verdera, F.[a], Pérez-Fernández, C.[b], Farias, M. E.[c], González, B.[d], Orfila, A.[e], Rosselló-Móra, R.[a]

[a] Department of Ecology and Marine Resources, Mediterranean Institute for Advanced Studies (IMEDEA, UIB-CSIC), Spain.

[b] Environmental Microbiology Laboratory, Puerto Rico University, Rio Piedras campus.

[c] Laboratorio de Investigaciones Microbiológicas de Lagunas Andinas (LIMLA), Planta Piloto de Procesos Industriales Microbiológicos (PROIMI), CCT, CONICET, San Miguel de Tucumán, Tucumán, Argentina.

[d] Facultad de Ingeniería y Ciencias, Universidad Adolfo Ibáñez – Center of Applied Ecology and Sustainability, Santiago, Chile.

[e] Marine Technology and Operational Oceanography Department, IMEDEA (CSIC-UIB), Esporles, Spain.

Corresponding author: Mora-Ruiz, Merit del R.

Marine Microbiology Group, Department of Ecology and Marine Resources, Mediterranean Institute for Advanced Studies, CSIC-UIB, C/Miquel Marqués 21, 07190 Esporles, Spain

Tel: +34 971 611 827 ; Email: meritmora@gmail.com

Abstract

Microorganisms are globally distributed but new evidence shows that the microbial structure of their communities can vary due to geographical location and environmental parameters. In this