Accepted Manuscript



A membrane disrupting toxin from wasp venom underlies the molecular mechanism of tissue damage

Rose Ombati, Yunfei Wang, Canwei Du, Xiancui Lu, Bowen Li, Atunga Nyachieo, Yaxiong Li, Shilong Yang, Ren Lai

PII: S0041-0101(18)30144-2

DOI: 10.1016/j.toxicon.2018.04.011

Reference: TOXCON 5860

To appear in: *Toxicon*

Received Date: 6 December 2017

Revised Date: 8 April 2018

Accepted Date: 10 April 2018

Please cite this article as: Ombati, R., Wang, Y., Du, C., Lu, X., Li, B., Nyachieo, A., Li, Y., Yang, S., Lai, R., A membrane disrupting toxin from wasp venom underlies the molecular mechanism of tissue damage, *Toxicon* (2018), doi: 10.1016/j.toxicon.2018.04.011.

A membrane disrupting toxin from wasp venom underlies the molecular mechanism of tissue damage

Rose Ombati[a,b,d]¶, Yunfei Wang[a,b]¶, Canwei Du[c], Xiancui Lu[a,b], Bowen Li[a,b], Atunga Nyachieo[d], Yaxiong Li[e]*, Shilong Yang[a]*, Ren Lai[a, c]*

[a] Key Laboratory of Animal Models and Human Disease Mechanisms of Chinese Academy of Sciences/Key Laboratory of bioactive peptides of Yunnan Province, Kunming Institute of Zoology, Kunming 650223, Yunnan, China.

[b] University of Chinese Academy of Sciences, Beijing 100049, China

[c] College of Life Sciences, Nanjing Agricultural University, Nanjing 210095, Jiangsu, China

[d] Institute of Primate Research, National Museums of Kenya, Nairobi, Kenya PO Box 24481-00502, Karen, Nairobi, Kenya.

[e] Cardiovascular Surgery Department of Kunming Yan'an Hospital, Kunming 650051, China

¶These authors contributed equally to this work.

*Correspondence and requests for materials should be addressed to Y.L. (liyaxiong62@aliyun.com), S.Y. (yangsl@mail.kiz.ac.cn) or to R.L. (rlai@mail.kiz.ac.cn).

Abstract

The molecular mechanism of the local hypersensitivity reactions to wasp venom including dermal necrosis remains an enigma regardless of the numerosity of the reported cases. In this study, we discovered a new membrane disrupting toxin, VESCP-M2 responsible for tissue