



Contents lists available at ScienceDirect

Toxicon

journal homepage: www.elsevier.com/locate/toxicon

Short communication

Beta-defensin genes of the Colubridae snakes *Phalotris mertensi*, *Thamnodynastes hypoconia*, and *T. strigatus*Yago Santana de Oliveira, Poliana G. Corrêa, Nancy Oguiura^{*}

Ecology and Evolution Laboratory – Instituto Butantan, Av. Dr. Vital Brasil 1500, 05503-900, São Paulo, Brazil

ARTICLE INFO

Article history:

Received 5 November 2017

Received in revised form

22 February 2018

Accepted 26 February 2018

Available online xxx

Keywords:

Antimicrobial peptides

Colubrid snakes

 β -defensin genes

ABSTRACT

β -Defensins are cationic antimicrobial peptides showing little sequence similarity but highly conserved tertiary structure stabilized by a six-cysteines-motif. Using a PCR approach, we described β -defensin sequences with two exons in three species of Colubridae snakes with high sequence similarity between them. The deduced amino acid sequence presented the characteristics of β -defensin family. The phylogenetic analysis using β -defensin coding sequences of different snakes grouped them in two main branches: genes organized in three or two exons.

© 2018 Elsevier Ltd. All rights reserved.

β -Defensins comprise a class of cationic antimicrobial peptides of innate immunity which perform functions in adaptive immunity and in non-immunological processes. β -Defensin family members have little sequence similarity, but a high degree of similarity in their tertiary structure: three β -strands arranged in an antiparallel sheet held together by the three intramolecular cysteine disulfide bonds: Cys1-Cys5, Cys2-Cys4 and Cys3-Cys6 (Machado and Ottoloni, 2015). The structure of pre- β -defensin consists of a signal sequence, a short or absent propeptide, and the mature defensin (Ganz, 2003). The first β -defensin described in snakes was crotamine, a small basic myotoxin from the venom of the rattlesnake *Crotalus durissus terrificus* (Oguiura et al., 2005; Coronado et al., 2013).

In reptiles, β -defensin-like genes have been described in snakes (Corrêa and Oguiura, 2013; Rádís-Baptista et al., 2004, 2003), and lizards (Dalla Valle et al., 2012) having three exons and two introns. In snake sequences, introns 1 and 2 are, respectively, of phase 1 (intron split a codon after the first nucleotide) and 2 (intron split a codon after the second nucleotide). The phase 1 intron after the signal peptide seems be usual to β -defensin genes (Zhu and Gao, 2013) and to proteins that will be exported (Sanz and Calvete, 2016).

The evolutionary relationship between defensins is still unclear

(Zhu and Gao, 2013); to understand the evolution of these genes in snakes and consequently in vertebrates, we analyzed β -defensin genes in the colubrid snakes *Phalotris mertensi*, *Thamnodynastes hypoconia* and *T. strigatus* using PCR, an approach similar to that used by Corrêa and Oguiura (2013), and Schutte and Mccray (2002).

The β -defensin genes of colubrid snakes were obtained by PCR amplification using primers designed based on cDNA sequences of Duvernoy glands from *T. strigatus* and *P. mertensi* (Ching et al., 2012; Campos et al., 2016). Genomic DNA was purified from liver obtained from the Tissue Bank of Herpetological Collection Romano Hoge of the Butantan Institute (*T. hypoconia*, Itú-SP, IBSP 83.377; *T. strigatus*, Itatiba-SP, IB-SP 83.628; and *P. mertensi*, Sumaré-SP, IBSP-82.259) using Chelex (Walsh et al., 1991). A 20- μ l reaction mix contained 60–500 ng DNA sample, 0.1 mM of each primer, 0.5 U Taq DNA Polymerase Platinum (Invitrogen), buffer with the addition of 2.5 mM MgCl₂, and 0.2 mM dNTP mix. The amplification process consisted of an initial denaturation step of 4 min at 94 °C, followed by 30 cycles of 45 s at 94 °C, 45 s at 58–65 °C and 45 s at 72 °C, and finally 1 min at 72 °C.

The amplicon was purified, after electrophoresis on a 1% agarose gel, using the Zymoclean Gel DNA Recovery kit (ZymoResearch). The purified DNA was cloned into the pTZ57 R/T vector according to the manufacturer's instructions (Fermentas). Ten microliters of ligation mixture were used to transform the *E. coli* DH5a (Ausubel et al., 2000). Clones were sequenced using the Sanger method and fractionated on an ABI Prism 3500 Genetic Analyzer (Applied Biosystems). Sequencing was performed at the Biotechnology Center in the Butantan Institute.

^{*} Corresponding author.

E-mail addresses: yagobiologia@gmail.com (Y.S. de Oliveira), poliana.correa@butantan.gov.br (P.G. Corrêa), nancy.oguiura@butantan.gov.br (N. Oguiura).

We analyzed seven sequences of *P. mertensi*, eight of *T. hypoconia*, and seven of *T. strigatus*. The sequences are showed in [Supplementary Material-1](#). These genes showed two exons. Like the other β -defensin genes of snakes and other animals, the exons 1 are interrupted after the first nucleotide of the last codon (phase 1 intron). The exon 1 codifies the signal peptide and exon 2 the last three amino acids of signal peptide and the mature peptide. Other snake β -defensin genes also codify the signal peptide in exon 1 and the last three amino acid of signal peptide and mature peptide in exons 2 and 3. Exons 1 were conserved while the second exon showed lower similarity between genes of *P. mertensi*, all sequences showed the six-cysteine motif, which is characteristic of β -defensins. The signal peptide was hydrophobic and leucine-rich. Signal P ([Petersen et al., 2011](#)) indicated that the cleavage site occurs after GNA and the first amino acid of mature peptide is Q in all β -defensins analyzed similar to crotoxin and other Brazilian pit vipers ([Corrêa and Oguiura, 2013](#)). The mature peptides had a predicted positive net charge at pH 7 and showed the consensus X₃₋₄-C-X₆-C-X₄-C-X₁₁-C-X₅-C-C-X₂ similar to that of other vertebrate β -defensins. Unlike the other described β -defensins of snakes, the carboxy-terminal tails of these colubrid snake defensins were short and those of *P. mertensi* were not positively charged ([Fig. 1](#)). The function of β -defensins of snakes are unknown but crotoxin has antimicrobial activities ([Yamane et al., 2013](#); [Oguiura et al., 2011](#); [Yount et al., 2009](#)) in addition to the myotoxic activity. These activities are due to its capacity to pore formation in artificial membrane of bacteria ([Costa et al., 2014](#)) and interaction with potassium channels ([Peigneur et al., 2012](#)). About the β -defensins studied here, we only know that synthetic linear peptides Defb_Ts and Defb_Bj-01 have some antimicrobial activity only against *Micrococcus luteus* (data not shown). It is necessary a specific investigation using a folded peptide in order to determine their function because the β -defensins can have multiple functions in innate immunity in addition to the antimicrobial activity ([Lai and Gallo, 2009](#)).

The seven *P. mertensi* genomic sequences coded for three different mature peptides but different to transcript previously described ([Campos et al., 2016](#)). Eight analyzed sequences of *T. hypoconia* coded for two sequences with one-amino acid difference and different compared to *T. strigatus*. Among the seven sequences of *T. strigatus*, six mature peptides were identical to the cDNA described before ([Ching et al., 2012](#)). We observed that Defb_Pm-06 and _Pm-07 showed a signal peptide identical to *Thamnodynastes* sequences, but diverse of one amino acid (E₁₈ instead Q₁₈) in relation to other *Phalotris* signal sequences. The greatest differences of both sequences occurred in mature peptides. On the other hand, *Thamnodynastes* sequences were more conserved in both regions. The variation between introns are mainly because of size variation ([Table 1](#)). The *Thamnodynastes*

Table 1

Sequence similarities of β -defensin genes of colubrid snake and *Bothrops jararaca* snake.

Sequence	GenBank	Exon 1	Intron 1	Exon 2	Intron 2	Exon 3
Defb_Pm-01	KX664436	—	—	—	—	—
Defb_Pm-03	KX664441	100%	99.5%	98.4%	—	—
Defb_Pm-04	KX664437	100%	92.7%	98.4%	—	—
Defb_Pm-05	KX664440	100%	99.3%	98.4%	—	—
Defb_Pm-06	KX664439	98.3%	85.7%	68.0%	—	—
Defb_Pm-07	KX664442	96.6%	93.0%	70.5%	—	—
Defb_Pm-08	KX664438	100%	99.5%	97.5%	—	—
Defb_Th-01	KX664421	—	—	—	—	—
Defb_Th-02	KX664422	100%	96.3%	100%	—	—
Defb_Th-03	KX664422	100%	99.5%	100%	—	—
Defb_Th-04	KX664424	100%	99.5%	99.2%	—	—
Defb_Th-05	KX664425	100%	99.5%	100%	—	—
Defb_Th-06	KX664426	100%	99.0%	100%	—	—
Defb_Th-07	KX664427	100%	89.0%	99.2%	—	—
Defb_Th-08	KX664428	100%	99.4%	100%	—	—
Defb_Ts-01	KX664429	—	—	—	—	—
Defb_Ts-05	KX664435	100%	86.2%	100%	—	—
Defb_Ts-06	KX664430	100%	85.9%	99.2%	—	—
Defb_Ts-07	KX664433	100%	86.4%	100%	—	—
Defb_Ts-08	KX664432	100%	83.9%	100%	—	—
Defb_Ts-10	KX664431	100%	85.9%	100%	—	—
Defb_Ts-21	KX664434	100%	86.3%	100%	—	—
Defb_Bj-04	MG833857	—	—	—	—	—
Defb_Bj-05	MG833858	100%	99.8%	100%	100%	100%
Defb_Bj-02	KC117164	100%	77.4%	76.3%	97.4%	93.8%
Defb_Bj-06	MG833859	100%	96.2%	100%	100%	100%
Defb_Bj-07	MG833860	100%	96.0%	100%	100%	100%
Defb_Bj-01	KC117163	100%	24.8%	80.5%	98.7%	75.0%
Defb_Bj-03	MG833861	100%	99.6%	100%	100%	100%

We used Geneious 6.0.6 ([Kearse et al., 2012](#)) to analyze the sequences based on Muscle alignment. Abbreviations are defined in [Fig. 1](#). Defb-Pm were compared to Defb-Pm01; Defb-Th to Defb-Th01; Defb-Ts to Defb-Ts01; and Defb_Bj to Defb_Bj-04. The sequences Defb_Bj of *B. jararaca* snake were obtained as described in [Corrêa and Oguiura \(2013\)](#). The similarity is presented as percentage of identity.

sequences seem to be more conserved than other defensin sequences of snakes.

To identify possible differences in the diversification patterns between the similarities of β -defensin genes with two or three exons, we also analyzed seven other genomic sequences of *Bothrops jararaca* ([Supplementary material-2](#)). We could observe a greater variation between the sizes of intron 1 and the mature peptide with the same signal peptide in pit viper sequences ([Table 1](#)).

Since the majority of β -defensin genes in mammals have two exons ([Patil et al., 2005](#)) and four exons in birds ([Xiao et al., 2004](#)), we supposed that in snakes the major structure would have three exons ([Corrêa and Oguiura, 2013](#)) as in lizards ([Dalla Valle et al., 2012](#)). Surprisingly, in the colubrid snake sequences analyzed herein, the β -defensin genes showed two exons, not three like the Brazilian pit viper snakes. This characteristic is not specific to




Fig. 1. Alignment of propeptide of β -defensin-coding sequences described in this work. Sequences were aligned using Muscle at Geneious 6.0.6 ([Kearse et al., 2012](#)) and the graphic view prepared using BioEdit ([Hall, 1999](#)). Defb_Pm, β -defensin genes of *Phalotris mertensi*; Defb_Th, β -defensin genes of *Thamnodynastes hypoconia*; Defb_Ts, β -defensin genes of *T. strigatus*. The green color indicates the polar amino acids, blue color the basic amino acids, red color the acidic amino acids, and brown color the cysteines. The predicted positive net charge at pH 7 and pI were calculated using PepDraw (<http://pepdraw.com/>) by Thomas C. Freeman, Jr.: Pm-07 (+3, 8.3); Pm-06 (+4, 8.5); Pm-03 (+6, 8.9); Ts-01 (+3, 8.3); Th-07 (+1, 7.6); Th-01 (+1, 7.6); Bj-02 (+9, 10.4); Bj-03 (+3, 8.2); Bj-01 (+4, 8.5).

Download English Version:

<https://daneshyari.com/en/article/8394567>

Download Persian Version:

<https://daneshyari.com/article/8394567>

[Daneshyari.com](https://daneshyari.com)