A genome-wide association study of mitochondrial DNA in Chinese men identifies two risk single nucleotide substitutions for idiopathic oligoasthenospermia

Chuncheng Lu, Miaofei Xu, Rong Wang, Yufeng Qin, Jing Ren, Wei Wu, Ling Song, Shoulin Wang, Zuomin Zhou, Hongbing Shen, Jiahao Sha, Zhibin Hu, Yankai Xia, Dengshun Miao, Xinru Wang

PII: S1567-7249(15)30014-3
DOI: doi: 10.1016/j.mito.2015.07.007
Reference: MITOCH 1024

To appear in: *Mitochondrion*

Received date: 25 September 2014
Revised date: 9 May 2015
Accepted date: 20 July 2015

Please cite this article as: Lu, Chuncheng, Xu, Miaofei, Wang, Rong, Qin, Yufeng, Ren, Jing, Wu, Wei, Song, Ling, Wang, Shoulin, Zhou, Zuomin, Shen, Hongbing, Sha, Jiahao, Hu, Zhibin, Xia, Yankai, Miao, Dengshun, Wang, Xinru, A genome-wide association study of mitochondrial DNA in Chinese men identifies two risk single nucleotide substitutions for idiopathic oligoasthenospermia, *Mitochondrion* (2015), doi: 10.1016/j.mito.2015.07.007

A genome-wide association study of mitochondrial DNA in Chinese men identifies two risk single nucleotide substitutions for idiopathic oligoasthenospermia

Chuncheng Lu[1, 2, †], Miaofei Xu[1, 2, †], Rong Wang[3, †], Yufeng Qin[1, 2], Jing Ren[3], Wei Wu[1, 2], Ling Song[1, 2], Shoulin Wang[1, 2], Zuomin Zhou[1], Hongbing Shen[1, 4], Jiahao Sha[1], Zhibin Hu[1, 4], Yankai Xia[1, 2], Dengshun Miao[1, 3, *] and Xinru Wang[1, 2, *]

[1] State Key Laboratory of Reproductive Medicine, Institute of Toxicology, Nanjing Medical University, Nanjing 210029, China

[2] Key Laboratory of Modern Toxicology of Ministry of Education, School of Public Health, Nanjing Medical University, Nanjing 210029, China

[3] Research Center for Bone and Stem Cells, Department of Anatomy, Histology, and Embryology, Nanjing Medical University, Nanjing, China.

[4] Department of Epidemiology and Biostatistics and Key Laboratory of Modern Toxicology of Ministry of Education, School of Public Health, Nanjing Medical University, Nanjing, China.

[†] The authors have contributed equally to this study and they should be regarded as joint first authors.

[*] To whom correspondence should be addressed at:

Drs. Dengshun Miao and Xinru Wang

State Key Laboratory of Reproductive Medicine, Institute of Toxicology, Nanjing Medical University, Nanjing 210029, China

Phone: +86-25-86862845; Fax: +86-25-86862847

E-mail: dengshun_miao@yahoo.com.cn and xrwang@njmu.edu.cn