


Author's Accepted Manuscript

N- and C-terminal regions of carrot Heat shock protein 17.7 can confer abiotic stress tolerance to transformed *Escherichia coli*

Eunhye Ko, Yeh-Jin Ahn



www.elsevier.com/locate/bab

PII: S1878-8181(18)30156-7
DOI: <https://doi.org/10.1016/j.bcab.2018.03.003>
Reference: BCAB715

To appear in: *Biocatalysis and Agricultural Biotechnology*

Received date: 13 February 2018
Revised date: 3 March 2018
Accepted date: 5 March 2018

Cite this article as: Eunhye Ko and Yeh-Jin Ahn, N- and C-terminal regions of carrot Heat shock protein 17.7 can confer abiotic stress tolerance to transformed *Escherichia coli*, *Biocatalysis and Agricultural Biotechnology*, <https://doi.org/10.1016/j.bcab.2018.03.003>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

N- and C-terminal regions of carrot Heat shock protein 17.7 can confer abiotic stress tolerance to transformed *Escherichia coli*

Eunhye Ko, Yeh-Jin Ahn*

Department of Biotechnology, Sangmyung University, 20 Hongjimun 2-gil, Jongno-gu, Seoul 03016, Korea

*Corresponding author.

E-mail address: yjahn@smu.ac.kr (Y. Ahn).

Keywords: abiotic stress; α -crystallin domain; *Daucus carota*; *Escherichia coli*; heat shock protein; molecular chaperone.

Download English Version:

<https://daneshyari.com/en/article/8405817>

Download Persian Version:

<https://daneshyari.com/article/8405817>

[Daneshyari.com](https://daneshyari.com)