

Accepted Manuscript

A predictive threshold value for the diagnosis of early pregnancy in cows using interferon-stimulated genes in granulocytes

Hitomi Yoshino, Noriyuki Toji, Kouya Sasaki, Katsuo Koshi, Norio Yamagishi, Toru Takahashi, Toshina Ishiguro-Oonuma, Hideo Matsuda, Tadayuki Yamanouchi, Yutaka Hashiyada, Kei Imai, Yoshiaki Izaike, Keiichiro Kizaki, Kazuyoshi Hashizume

PII: S0093-691X(17)30555-1

DOI: [10.1016/j.theriogenology.2017.11.014](https://doi.org/10.1016/j.theriogenology.2017.11.014)

Reference: THE 14346

To appear in: *Theriogenology*

Received Date: 27 July 2017

Revised Date: 10 November 2017

Accepted Date: 11 November 2017

Please cite this article as: Yoshino H, Toji N, Sasaki K, Koshi K, Yamagishi N, Takahashi T, Ishiguro-Oonuma T, Matsuda H, Yamanouchi T, Hashiyada Y, Imai K, Izaike Y, Kizaki K, Hashizume K, A predictive threshold value for the diagnosis of early pregnancy in cows using interferon-stimulated genes in granulocytes, *Theriogenology* (2017), doi: 10.1016/j.theriogenology.2017.11.014.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1
2
3 A predictive threshold value for the diagnosis of early pregnancy in cows using interferon-stimulated
4 genes in granulocytes

5
6 Hitomi Yoshino ^{a,*}, Noriyuki Toji ^{a,b}, Kouya Sasaki ^c, Katsuo Koshi ^{a,b,**}, Norio Yamagishi ^{a,b,***},
7 Toru Takahashi ^{a,b}, Toshina Ishiguro-Oonuma ^{a,b}, Hideo Matsuda ^d, Tadayuki Yamanouchi ^d, Yutaka
8 Hashiyada ^d, Kei Imai ^e, Yoshiaki Izaike ^{a,b}, Keiichiro Kizaki ^{a,b}, Kazuyoshi Hashizume ^{a,b,****}

9
10
11 ^a Laboratory of Veterinary Physiology, Cooperative Department of Veterinary Science, Faculty of
12 Agriculture, Iwate University, 3-18-8 Ueda, Morioka, Iwate 020-8550, Japan

13 ^b United Graduate School of Veterinary Sciences, Gifu University, Gifu 501-1193, Japan

14 ^c Iwate Veterinary Hospital, Itsukaichi, Iwate Town, Iwate 028-4307, Japan

15 ^d National Livestock Breeding Center, Nishigo, Fukushima 961-8061, Japan

16 ^e Department of Sustainable Agriculture, Rakuno Gakuen University, 582 Bunkiyodai-Midorimachi,
17 Ebetsu, Hokkaido 069-8501, Japan

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32 * Present address: Hokkaido Research Organization, Agricultural Research Department Animal
33 Research Center, Hokkaido 081-0038, Japan

34 ** Present address: Gifu Prefectural Seino Public Health Center, Gifu 503-0838, Japan

35 *** Present address: Obihiro University, Hokkaido 080-8555, Japan

36 **** Corresponding author. Laboratory of Veterinary Physiology, Department of Veterinary
37 Medicine, Faculty of Agriculture, Iwate University, 3-18-8 Ueda, Morioka, Iwate 020-8550, Japan.
38 Tel./fax: +81 19 621 6210.

39 E-mail address: kazuha@iwate-u.ac.jp (K. Hashizume).

Download English Version:

<https://daneshyari.com/en/article/8427749>

Download Persian Version:

<https://daneshyari.com/article/8427749>

[Daneshyari.com](https://daneshyari.com)