


# Accepted Manuscript

Triploid or hybrid tetra: Which is the ideal sterile host for surrogate technology?

Lucas Henrique Piva, Diógenes Henrique de Siqueira-Silva, Caio Augusto Gomes Goes, Takafumi Fujimoto, Taiju Saito, Leticia Veroni Dragone, José Augusto Senhorini, Fabio Porto-Foresti, José Bento Sterman Ferraz, George Shigueki Yasui



PII: S0093-691X(17)30602-7

DOI: [10.1016/j.theriogenology.2017.12.013](https://doi.org/10.1016/j.theriogenology.2017.12.013)

Reference: THE 14383

To appear in: *Theriogenology*

Received Date: 16 May 2017

Revised Date: 2 December 2017

Accepted Date: 3 December 2017

Please cite this article as: Piva LH, de Siqueira-Silva DióHenrique, Goes CAG, Fujimoto T, Saito T, Dragone LetíVeroni, Senhorini JoséAugusto, Porto-Foresti F, Ferraz JoséBentoSterman, Yasui GS, Triploid or hybrid tetra: Which is the ideal sterile host for surrogate technology?, *Theriogenology* (2018), doi: 10.1016/j.theriogenology.2017.12.013.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Triploid or hybrid tetra: which is the ideal sterile host for surrogate technology?**

5 **Lucas Henrique Piva<sup>1,2</sup>, Diógenes Henrique de Siqueira-Silva<sup>2,3,4\*</sup>, Caio Augusto Gomes Goes<sup>5</sup>, Takafumi Fujimoto<sup>6</sup>, Taiju Saito<sup>7</sup>, Letícia Veroni Dragone<sup>2</sup>, José Augusto Senhorini<sup>2</sup>, Fabio Porto-Foresti<sup>5</sup>, José Bento Sterman Ferraz<sup>4</sup>, George Shigueki Yasui<sup>2,4\*</sup>**

10 <sup>1</sup>*UNESP – Univ. Estadual Paulista, Campus de Botucatu, Programa de Pós-Graduação em Ciências Biológicas (Zoologia) Botucatu, São Paulo, Brazil.*

<sup>2</sup>*Centro Nacional de Pesquisa e Conservação da Biota Aquática Continental (CEPTA-ICMBIO), Pirassununga, São Paulo, Brazil.*

15 <sup>3</sup>*UNIFESSPA – Universidade Federal do Sul e Sudeste do Pará. Instituto de Estudo em Saúde e Biológicas (IESB) Marabá, Pará, Brazil.*

<sup>4</sup>*USP– University of São Paulo, Faculdade de Zootecnia e Engenharia de Alimentos Departamento de Medicina Veterinária, Pirassununga, São Paulo, Brazil.*

<sup>5</sup>*UNESP – Univ. Estadual Paulista, Campus de Bauru, Faculdade de Ciências Bauru, São Paulo, Brazil.*

20 <sup>6</sup>*Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho, 041-8611 Hakodate, Japan*

<sup>7</sup>*Nishiura Station, South Ehime Fisheries Research Center, Ehime University, Uchidomari, Ainan, Japan.*

25

\* Supported by FAPESP 2010/17429-1 and 2011/11664-1

\*Correspondence: Diógenes Henrique de Siqueira-Silva - UNIFESSPA – Universidade Federal do Sul e Sudeste do Pará. Instituto de Estudo em Saúde e Biológicas (IESB) Marabá, Pará, Brazil. Folha 31, Quadra 07, Lote especial s/n Nova Marabá - Marabá/PA

30 CEP: 68.507-590. Email: [diogenessilva@unifesspa.edu.br](mailto:diogenessilva@unifesspa.edu.br); [siqueira.diogenes@gmail.com](mailto:siqueira.diogenes@gmail.com)

Download English Version:

<https://daneshyari.com/en/article/8427789>

Download Persian Version:

<https://daneshyari.com/article/8427789>

[Daneshyari.com](https://daneshyari.com)