

Uniqueness of the positive solution for a class of semilinear elliptic systems

Renhao Cui^a, Yuwen Wang^a, Junping Shi^{b,c,*}

^a *Y.Y. Tseng's Functional Research Center, Department of Mathematics, Harbin Normal University, Harbin, Heilongjiang 150025, PR China*

^b *Department of Mathematics, College of William and Mary, Williamsburg, VA 23187, USA*

^c *Department of Mathematics, Harbin Normal University, Harbin, Heilongjiang 150025, PR China*

Received 15 May 2006; accepted 16 August 2006

Abstract

We prove the uniqueness of the positive radially symmetric solution to the following problem:

$$\begin{cases} \Delta u + \lambda v^{p_1} = 0, & x \in B_1, \\ \Delta v + \lambda w^{p_2} = 0, & x \in B_1, \\ \Delta w + \lambda u^{p_3} = 0, & x \in B_1, \\ u = v = w = 0, & x \in \partial B_1, \end{cases}$$

where $p_i > 0$ ($i = 1, 2, 3$) and B_1 is the unit ball in \mathcal{R}^n .

© 2006 Elsevier Ltd. All rights reserved.

MSC: 35J55; 35B32

Keywords: Semilinear elliptic systems; Positive solution; Uniqueness

1. Introduction

In this paper we study the positive radially symmetric solutions of the semilinear elliptic system

$$\begin{cases} \Delta u + \lambda v^{p_1} = 0, & x \in B_1, \\ \Delta v + \lambda w^{p_2} = 0, & x \in B_1, \\ \Delta w + \lambda u^{p_3} = 0, & x \in B_1, \\ u = v = w = 0, & x \in \partial B_1, \end{cases} \quad (1.1)$$

where B_1 is the unit ball in \mathcal{R}^n , $n \geq 1$, and $p_i > 0$ ($i = 1, 2, 3$). By a transformation $U(y) = u(\lambda^{-\frac{1}{2}}y)$, $V(y) = v(\lambda^{-\frac{1}{2}}y)$, $W(y) = w(\lambda^{-\frac{1}{2}}y)$, we can convert (1.1) to

* Corresponding author at: Department of Mathematics, College of William and Mary, Williamsburg, VA 23187, USA.
E-mail addresses: renhaocui@163.com (R. Cui), wangyuwen2003@sohu.com (Y. Wang), shij@math.wm.edu (J. Shi).

$$\begin{cases} \Delta U + V^{p_1} = 0, & y \in B_R, \\ \Delta V + W^{p_2} = 0, & y \in B_R, \\ \Delta W + U^{p_3} = 0, & y \in B_R, \\ U = V = W = 0, & y \in \partial B_R. \end{cases} \tag{1.2}$$

We shall study (1.1) instead of (1.2) since the structure of the solution set of (1.2) is same as that of (1.1). Our approach to the uniqueness is based on two ingredients: (a) the parameterization of the set of all solutions; and (b) the scaling of the homogeneous equation (1.1). To illustrate the ideas, we consider the positive radial solutions of a scalar equation:

$$\begin{cases} \Delta u + \lambda u^p = 0, & x \in B_1, \\ u = 0, & x \in \partial B_1. \end{cases} \tag{1.3}$$

Like for (1.2), the solutions of (1.3) are equivalent to those of

$$\begin{cases} \Delta U + U^p = 0, & y \in B_R, \\ U = 0, & y \in \partial B_R, \end{cases} \tag{1.4}$$

via the same change of variables as above. Then from the uniqueness of the initial value problem of the ordinary differential equation, the radius R in (1.4) is uniquely determined by $U(0) = \max_{y \in \overline{B_R}} U(y)$, and so is $\lambda = R^2$. Thus the solution set of (1.4) is parameterized by a single parameter $U(0)$. On the other hand, if $u_1(x)$ is a solution of (1.3) with $\lambda = 1$, then $u_\lambda(x) = \lambda^{1/(1-p)}u_1(x)$ is a solution of (1.3) for general $\lambda > 0$, and the range of $\{u_\lambda(0)\}$ is \mathcal{R}^+ . The curve $\Sigma = \{(\lambda, u_\lambda) : \lambda > 0\}$ is monotone; hence we obtain the uniqueness of the solution for each $\lambda > 0$.

We follow a similar approach for the uniqueness of solution to the system (1.1). We generalize an idea of Dalmasso [2] and Korman and Shi [3] to prove that the solution set $\{(\lambda, u, v, w)\}$ of (1.1) (or equivalently $\{(R, U, V, W)\}$ of (1.2)) can be parameterized by a single variable u_0 (or U_0 respectively) under certain conditions. In particular, we prove the uniqueness of the solution of (1.1) for any fixed λ when $p_i > 0$ ($i = 1, 2, 3$), which generalizes results of [1,2,4]. In [5], the uniqueness of positive solution is obtained when the exponents are sublinear.

We state our main result.

Theorem 1.1. *We assume $p_i > 0$ ($i = 1, 2, 3$) and there exists $\lambda_0 > 0$ such that (1.1) has a positive radially symmetric solution $(u_{\lambda_0}, v_{\lambda_0}, w_{\lambda_0})$. Then:*

1. *If $1 - p_1 p_2 p_3 \neq 0$, then for each $\lambda > 0$, there exists exactly one positive radially symmetric solution $(u_\lambda, v_\lambda, w_\lambda)$.*
2. *If $1 - p_1 p_2 p_3 = 0$, then (1.1) has no positive radially symmetric solution for any $\lambda > 0$ and $\lambda \neq \lambda_0$, and (1.1) has infinitely many positive radially symmetric solutions at $\lambda = \lambda_0$, which can be represented as*

$$\left\{ \left(k u_{\lambda_0}, k^{\frac{1+p_2+p_2 p_3}{1+p_1+p_1 p_2}} v_{\lambda_0}, k^{\frac{1+p_3+p_1 p_3}{1+p_1+p_1 p_2}} w_{\lambda_0} \right) : k > 0 \right\}.$$

For (1.1), we call the system sublinear when $1 - p_1 p_2 p_3 > 0$, and we call it superlinear when $1 - p_1 p_2 p_3 < 0$. We prove the uniqueness of the solution in both sublinear and superlinear cases.

Our uniqueness result is proved under the assumption that a positive solution exists. The existence results for the system in (1.1) on general domains for the exponents satisfying $1 - p_1 p_2 p_3 > 0$ have been proved in [5] (see Theorem 1.2 below), but the existence for the superlinear case is still not known.

Theorem 1.2. *Consider the problem*

$$\begin{cases} \Delta u + v^{p_1} = 0, & x \in D, \\ \Delta v + w^{p_2} = 0, & x \in D, \\ \Delta w + u^{p_3} = 0, & x \in D, \\ u = v = w = 0, & x \in \partial D, \end{cases} \tag{1.5}$$

where D is a ball in \mathcal{R}^n , $p_i > 0$ ($i = 1, 2, 3$) and $1 - p_1 p_2 p_3 > 0$. Then (1.5) admits a positive classical solution (u, v, w) .

Download English Version:

<https://daneshyari.com/en/article/843190>

Download Persian Version:

<https://daneshyari.com/article/843190>

[Daneshyari.com](https://daneshyari.com)