

## Accepted Manuscript

Nuclear BAP1 loss is common in intrahepatic cholangiocarcinoma and a subtype of hepatocellular carcinoma but rare in pancreatic ductal adenocarcinoma

Asmaa Mosbeh , Khalil Halfawy , Wael S. Abdel-Mageed ,  
Dina Sweed , Mohamed H. Abdel Rahman

PII: S2210-7762(17)30357-5  
DOI: [10.1016/j.cancergen.2018.03.002](https://doi.org/10.1016/j.cancergen.2018.03.002)  
Reference: CGEN 587

To appear in: *Cancer Genetics*

Received date: 15 September 2017  
Revised date: 17 January 2018  
Accepted date: 29 March 2018

Please cite this article as: Asmaa Mosbeh , Khalil Halfawy , Wael S. Abdel-Mageed , Dina Sweed , Mohamed H. Abdel Rahman , Nuclear BAP1 loss is common in intrahepatic cholangiocarcinoma and a subtype of hepatocellular carcinoma but rare in pancreatic ductal adenocarcinoma, *Cancer Genetics* (2018), doi: [10.1016/j.cancergen.2018.03.002](https://doi.org/10.1016/j.cancergen.2018.03.002)



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights**

- Loss of BAP1 is common in cholangiocarcinoma and a subtype of hepatocellular carcinoma but rare in pancreatic cancer.
- Hepatocellular carcinomas with BAP1 loss show co-expression of biliary and hepatic epithelial markers suggesting that they are a unique subtype of the tumor.
- Loss of heterozygosity of the 3p21 region is commonly detected in the hepatobiliary and pancreatic tumors.

Download English Version:

<https://daneshyari.com/en/article/8433521>

Download Persian Version:

<https://daneshyari.com/article/8433521>

[Daneshyari.com](https://daneshyari.com)