

Accepted Manuscript

Mutation analysis of therapy-related myeloid neoplasms

Takahiro Nishiyama , Yuichi Ishikawa , Naomi Kawashima ,
Akimi Akashi , Yoshiya Adachi , Hikaru Hattori , Yoko Ushijima ,
Hitoshi Kiyoi

PII: S2210-7762(18)30004-8
DOI: [10.1016/j.cancergen.2018.02.006](https://doi.org/10.1016/j.cancergen.2018.02.006)
Reference: CGEN 583

To appear in: *Cancer Genetics*

Received date: 5 January 2018
Revised date: 17 February 2018
Accepted date: 21 February 2018

Please cite this article as: Takahiro Nishiyama , Yuichi Ishikawa , Naomi Kawashima , Akimi Akashi , Yoshiya Adachi , Hikaru Hattori , Yoko Ushijima , Hitoshi Kiyoi , Mutation analysis of therapy-related myeloid neoplasms, *Cancer Genetics* (2018), doi: [10.1016/j.cancergen.2018.02.006](https://doi.org/10.1016/j.cancergen.2018.02.006)



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- *t*-MN cells preferentially acquired *TP53* and epigenetic modifying genes mutations.
- A somatic *TET2* mutation was identified in the normal PB at the disease-free period.
- A germ-line *MYB* mutation was identified in a patient who developed MDS after FL.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/8433654>

Download Persian Version:

<https://daneshyari.com/article/8433654>

[Daneshyari.com](https://daneshyari.com)