

Genetically-modified bone mesenchymal stem cells with TGF- β_3 improve wound healing and reduce scar tissue formation in a rabbit model

Mingyong Li, Lin Qiu, Wei Hu, Xiang Deng, Hanfeng Xu, Youhan Cao, Zijian Xiao, Liangyu Peng, Shawn Johnson, Lyundup Alexey, Paul A. Kingston, Qing Li, Yuanyuan Zhang



PII: S0014-4827(18)30080-6
DOI: <https://doi.org/10.1016/j.yexcr.2018.02.006>
Reference: YEXCR10919

To appear in: *Experimental Cell Research*

Received date: 31 August 2017
Revised date: 11 February 2018
Accepted date: 13 February 2018

Cite this article as: Mingyong Li, Lin Qiu, Wei Hu, Xiang Deng, Hanfeng Xu, Youhan Cao, Zijian Xiao, Liangyu Peng, Shawn Johnson, Lyundup Alexey, Paul A. Kingston, Qing Li and Yuanyuan Zhang, Genetically-modified bone mesenchymal stem cells with TGF- β_3 improve wound healing and reduce scar tissue formation in a rabbit model, *Experimental Cell Research*, <https://doi.org/10.1016/j.yexcr.2018.02.006>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Genetically-modified bone mesenchymal stem cells with TGF- β_3 improve wound healing and reduce scar tissue formation in a rabbit model

Mingyong Li¹, Lin Qiu², Wei Hu^{1*}, Xiang Deng¹, Hanfeng Xu¹, Youhan Cao¹, Zijian Xiao¹, Liangyu Peng¹, Shawn Johnson³, Lyundup Alexey⁴, Paul A. Kingston⁵, Qing Li^{1*}, Yuanyuan Zhang^{3*}

¹First affiliated hospital, University of South China, Hengyang, Hunan, China.

²Children's Hospital of Chongqing Medical University, Chongqing, China.

³Wake Forest Institute for Regenerative Medicine, Wake Forest University, North Carolina, USA.

⁴Advanced Cell Technologies Department of Institute for Regenerative Medicine and Regenerative Medicine Facility Center of Sechenov First Moscow State Medical University, Russia.

⁵Vascular Gene Therapy Unit, University of Manchester, Manchester, UK.

Huwei_abc970642@163.com

liq73@163.com

yzhang@wakehealth.edu;

*** Correspondence to:** Telephone: +86 180 0734 4154; Fax: +86 0734 8578592

*** Correspondence to:** No. 69, chuanshan Road, Shigu District, Hengyang 421001, China

*** Correspondence to:** Telephone: +86-136-1121-8973. Fax: +86-02363631928.

Abstract

Extensive scar tissue formation often occurs after severe burn injury, trauma, or as one of complications after surgical intervention. Despite significant therapeutic advances, it is still a

Download English Version:

<https://daneshyari.com/en/article/8450721>

Download Persian Version:

<https://daneshyari.com/article/8450721>

[Daneshyari.com](https://daneshyari.com)