www.elsevier.com/locate/yexcr

Exosomes derived from imatinib-resistant chronic myeloid leukemia cells mediate a horizontal transfer of drug-resistant trait by delivering miR-365

Qing-Hua Min, Xiao-Zhong Wang, Jing Zhang, Qing-Gen Chen, Shu-Qi Li, Xiao-Qing Liu, Jing Li, Jing Liu, Wei-Ming Yang, Yu-Huan Jiang, Yan-Mei Xu, Jin Lin, Qiu-Fang Gao, Fan Sun, Lei Zhang, Bo Huang

PII: S0014-4827(17)30643-2
DOI: https://doi.org/10.1016/j.yexcr.2017.12.001
Reference: YEXCR10840

To appear in: *Experimental Cell Research*

Received date: 6 April 2017
Revised date: 27 November 2017
Accepted date: 3 December 2017

Cite this article as: Qing-Hua Min, Xiao-Zhong Wang, Jing Zhang, Qing-Gen Chen, Shu-Qi Li, Xiao-Qing Liu, Jing Li, Jing Liu, Wei-Ming Yang, Yu-Huan Jiang, Yan-Mei Xu, Jin Lin, Qiu-Fang Gao, Fan Sun, Lei Zhang and Bo Huang, Exosomes derived from imatinib-resistant chronic myeloid leukemia cells mediate a horizontal transfer of drug-resistant trait by delivering miR-365, *Experimental Cell Research,* https://doi.org/10.1016/j.yexcr.2017.12.001

Exosomes derived from imatinib-resistant chronic myeloid leukemia cells mediate a horizontal transfer of drug-resistant trait by delivering miR-365

Qing-Hua Min[1], Xiao-Zhong Wang[1], Jing Zhang[1], Qing-Gen Chen[1], Shu-Qi Li[1], Xiao-Qing Liu[2], Jing Li[3], Jing Liu[1], Wei-Ming Yang[1],Yu-Huan Jiang[1], Yan-Mei Xu[1], Jin Lin[1], Qiu-Fang Gao[1], Fan Sun[1], Lei Zhang[1], Bo Huang[1]*

1 Department of Clinical Laboratory, the Second Affiliated Hospital of Nanchang University, Nanchang, China

2 The Medical School of Nanchang University, Nanchang, China

3 Departments of Clinical Laboratory, the First Affiliated Hospital of Nanchang University, Nanchang, China

*Corresponding author: Bo Huang, Department of Clinical Laboratory, the Second Affiliated Hospital of Nanchang University, No.1 Min De Road, Nanchang 330006, China E-mail: 764019522@qq.com Tel: 0086-791-86300410 Fax: 0086-791-86262262

Qing-Hua Min and Xiao-Zhong Wang are joint first authors.

Abstract