Accepted Manuscript

Title: Log odds of positive lymph nodes may predict survival benefit in patients with node-positive non-small cell lung cancer

Authors: Weiye Deng, Ting Xu, Yifan Wang, Yujin Xu, Pei Yang, Daniel Gomez, Zhongxing Liao



PII:	S0169-5002(18)30381-7
DOI:	https://doi.org/10.1016/j.lungcan.2018.05.016
Reference:	LUNG 5653
To appear in:	Lung Cancer
Received date:	6-2-2018
Revised date:	2-5-2018
Accepted date:	19-5-2018

Please cite this article as: Deng Weiye, Xu Ting, Wang Yifan, Xu Yujin, Yang Pei, Gomez Daniel, Liao Zhongxing.Log odds of positive lymph nodes may predict survival benefit in patients with node-positive non-small cell lung cancer.*Lung Cancer* (2018), https://doi.org/10.1016/j.lungcan.2018.05.016

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Log odds of positive lymph nodes may predict survival benefit in patients with nodepositive non-small cell lung cancer

Weiye Deng, MD, MPH¹, Ting Xu, MD, PhD¹, Yifan Wang, BS², Yujin Xu, MD³, Pei Yang, MD, MPH⁴, Daniel Gomez, MD¹, Zhongxing Liao, MD¹.

¹Department of Radiation Oncology, The University of Texas MD Anderson Cancer Center, 1400 Pressler St., Houston, TX 77030-4008; ²Department of Experimental Radiation Oncology, The University of Texas MD Anderson Cancer Cancer, 1515 Holcombe Blvd., Houston, TX 77030; ³Department of Radiation Oncology, Zhejiang Cancer Hospital, N0. 1, Banshandong Road, Gongshu District, Hanzhou, 310022, China; ⁴Department of 3rd Radiation Oncology, Hunan Cancer Hospital, 283 Tongzipo Road, Yuelu District, Changsha, Hunana, 410013, China.

Corresponding author: Zhongxing Liao, MD, Department of Radiation Oncology, Unit 1422, The University of Texas MD Anderson Cancer Center, 1400 Pressler St., Houston, TX 77030-4008. Tel 713-563-2300; E-mail: zliao@mdanderson.org.

Funding: None.

Word count of the manuscript text: 3147.

Highlights:

- □ LODDS, npLNs, and LNR were independent risk factors for OS and CSS.
- □ Higher LODDS was associated with worse OS and worse CSS.
- □ A LODDS model had the best fit compared with LNR or npLN models in predicting OS and CSS.
- □ LODDS and LNR staging schemes outperformed those of npLNs for predicting OS and CSS.

Download English Version:

https://daneshyari.com/en/article/8453664

Download Persian Version:

https://daneshyari.com/article/8453664

Daneshyari.com