

## Accepted Manuscript

Title: Morphological features of the anterior talofibular ligament by the number of fiber bundles

Authors: Mutsuaki Edama, Ikuo Kageyama, Takanori Kikumoto, Masatoshi Nakamura, Wataru Ito, Emi Nakamura, Ryo Hirabayashi, Tomoya Takabayashi, Takuma Inai, Hideaki Onishi



PII: S0940-9602(17)30147-4  
DOI: <https://doi.org/10.1016/j.aanat.2017.11.001>  
Reference: AANAT 51200

To appear in:

Received date: 13-7-2017  
Revised date: 6-10-2017  
Accepted date: 8-11-2017

Please cite this article as: Edama, Mutsuaki, Kageyama, Ikuo, Kikumoto, Takanori, Nakamura, Masatoshi, Ito, Wataru, Nakamura, Emi, Hirabayashi, Ryo, Takabayashi, Tomoya, Inai, Takuma, Onishi, Hideaki, Morphological features of the anterior talofibular ligament by the number of fiber bundles. *Annals of Anatomy* <https://doi.org/10.1016/j.aanat.2017.11.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **Morphological features of the anterior talofibular ligament by the number of fiber bundles**

Running headline: Classification by number of fiber bundles

Mutsuaki Edama<sup>1,2</sup>, Ikuo Kageyama<sup>2</sup>, Takanori Kikumoto<sup>1</sup>, Masatoshi Nakamura<sup>1</sup>, Wataru Ito<sup>1</sup>, Emi Nakamura<sup>1</sup>, Ryo Hirabayashi<sup>1</sup>, Tomoya Takabayashi<sup>1</sup>, Takuma Inai<sup>1</sup>, Hideaki Onishi<sup>1</sup>

<sup>1</sup> Institute for Human Movement and Medical Sciences, Niigata University of Health and Welfare, Shimami-cho 1398, Kita-ku, Niigata City, Niigata 950-3198, Japan

<sup>2</sup> Department of Anatomy, School of Life Dentistry at Niigata, Nippon Dental University, Niigata 951-8580, Japan

Corresponding Author: Mutsuaki Edama, Institute for Human Movement and Medical Sciences, Niigata University of Health and Welfare, Shimami-cho 1398, Kita-ku, Niigata City 950-3198, Japan, Tel: +81 25 257 4723, Fax: +81 25 257 4723, E-mail: edama@nuhw.ac.jp

### **Abstract**

The aims of this study have been to clarify differences in morphological features based on the number of fiber bundles in the anterior talofibular ligament (ATFL), and to investigate the relationship between the ATFL and the calcaneofibular ligament (CFL).

This study used 81 legs from 43 cadavers. The ATFL was classified according to

Download English Version:

<https://daneshyari.com/en/article/8460336>

Download Persian Version:

<https://daneshyari.com/article/8460336>

[Daneshyari.com](https://daneshyari.com)