


HOSTED BY



Contents lists available at ScienceDirect

The Egyptian Journal of Medical Human Genetics

journal homepage: www.sciencedirect.com

Original article

Evaluation of the association of single nucleotide polymorphisms in DDP4 and CDK5RAP2 genes with rheumatoid arthritis susceptibility in Iranian population

Zahra Malekshahi^{a,b}, Mahdi Mahmoudi^{a,*}, Massoomeh Akhlaghi^a, Masoud Garshasbi^c, Ahmadreza Jamshidi^a, Shiva Poursani^a, Shayan Mostafaei^a, Mohammad Hossein Nicknam^{b,*}

^a Rheumatology Research Center, Tehran University of Medical Sciences, Tehran, Iran

^b Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

^c Department of Medical Genetics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

ARTICLE INFO

Article history:

Received 17 May 2017

Accepted 7 June 2017

Available online xxxx

Keywords:

Rheumatoid arthritis

Gene polymorphism

Inflammation

Autoimmunity

DPP4

CDK5RAP2

ABSTRACT

Background: Rheumatoid arthritis (RA) is known as a chronic autoimmune inflammatory disorder, which is characterized mainly by the progressive inflammation and destruction of the joints. In the pathogenesis of RA, a variety of cell types such as lymphocyte, dendritic cells, osteoclasts and synovial fibroblasts are involved. Genetic proneness has been implicated in the pathogenesis of RA. The aim of this study was to evaluate the association of single nucleotide polymorphisms (SNPs) in *DPP4* and *CDK5RAP2* genes and risk of RA in Iranian population.

Methods: For genotyping, 623 RA patients and 412 healthy subjects were recruited. Genetic analysis of *DPP4* gene rs12617656 and *CDK5RAP2* gene rs12379034 polymorphisms was conducted using TaqMan allelic discrimination (for rs12617656) and ARMS-PCR (for rs12379034) methods.

Results: Experiments demonstrated that alleles and genotypes of both SNPs were represented equally in RA patients and controls. Statistical analysis revealed that none of the rs12617656 and rs12379034 SNPs had significant differences in prevalence of both alleles and genotypes between RA patients and healthy controls.

Conclusions: It appears that gene polymorphisms of *DPP4* and *CDK5RAP2* are not involved in the pathogenesis of RA in Iranian population.

© 2017 Ain Shams University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Abbreviations: RA, rheumatoid arthritis; SNP, single nucleotide polymorphism; GWAS, genome-wide association study; HLA, human leukocyte antigen; DPP4, dipeptidyl peptidase-4; CD26, cluster of differentiation 26; CDK-5, Cyclin-dependent kinase 5; CDK5RAP2, Cyclin-dependent kinase 5 regulatory subunit-associated protein 2; ACR, American college of rheumatology; EDTA, ethylenediaminetetraacetic acid; ARMS-PCR, amplification refractory mutation system-polymerase chain reaction; OR, odds ratio; HWE, Hardy-Weinberg equilibrium; ESR, erythrocyte sedimentation rate; DAS28, disease activity score in 28 joints; RF, rheumatoid factor; anti-CCP, anti-cyclic citrullinated peptide antibody; RANK, receptor activator of nuclear factor κ B.

Peer review under responsibility of Ain Shams University.

* Corresponding authors at: Rheumatology Research Center, Shariati Hospital, Kargar Ave., Tehran, Iran (M. Mahmoudi). Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Poursina Ave, 1417613151 Tehran, Iran (M.H. Nicknam).

E-mail addresses: mahmoudim@tums.ac.ir (M. Mahmoudi), nicknam_m@yahoo.com (M.H. Nicknam).

<http://dx.doi.org/10.1016/j.ejmhg.2017.06.002>

1110-8630/© 2017 Ain Shams University. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

RA is a chronic multifactorial disorder with involvement of the small diarthrodial joints of the hands and feet. Both genetics and environmental factors are the important contributing factors in the pathogenesis of RA [1–4]. Recently, determining roles of genetics in RA susceptibility have been explored mostly by genome-wide association studies (GWAS). Human leukocyte antigen (HLA) account for one-third of the genetic contributing factors for RA pathogenesis in addition to non-HLA components, which are increasingly being detected by genome-wide scanings [5].

DPP4 gene encodes dipeptidyl peptidase-4 (DPP4) or cluster of differentiation 26 (CD26), which is a membranous glycoprotein and is expressed on the surface of immune and non-immune cells with enzymatic activity of cleaving proline containing dipeptides. Its role has been clarified in some previous studies as an immune regulator and its serum levels and cell surface form have been sur-

veyed in several autoimmune diseases including RA [6,7]. The strong association of *DPP4* gene rs12617656 SNP with RA risk was demonstrated in a GWAS, indicating how genes can affect RA proneness in Han Chinese populations [8]. We also hypothesized that this polymorphism might take part in RA susceptibility and possibly in the progression of the disease in Iranian population.

Cyclin-dependent kinase 5 (CDK-5) regulatory subunit-associated protein 2 (*CDK5RAP2*) has an important role in differentiation process of neuronal system by forming the microtubule nucleator via gamma-tubulin ring complex recruitment to centrosomes [9]. This is a vital stage in brain evolution and mutations in the attributed genes during brain development result in mitosis division disturbance in neurons, culminating in microcephaly [10]. Surprisingly, a GWAS detected a strong association of rs12379034 SNP in *CDK5RAP2* gene with the above-mentioned function with RA susceptibility, introducing the association of a neuronal-related gene with an autoimmune disease for the first time [8].

Given the important roles of the *DPP4* gene in RA predisposition and how *CDK5RAP2* acts in immune and neuronal systems of RA patients, we were interested in surveying the possible connections between the two mentioned polymorphisms in these genes and RA risk in Iranian population.

2. Subjects and methods

2.1. Patients and controls

Our study population was composed of 623 (524 females and 99 male) RA patients recruited from the Rheumatology Clinic of Shariati Hospital and 412 (356 females and 56 male) healthy unrelated subjects without any history of autoimmune diseases. The RA patients who met the 1997 revised American College of Rheumatology (ACR) classification criteria [11] were delicately matched with control group considering age, sex, and ethnicity. Alongside genotyping of SNPs, phenotypical characteristics of RA patients were obtained to investigate the relation between two polymorphisms and clinical features. Written informed consent was taken by all the subjects and the protocol of this study was approved by the Ethics Committee of Tehran University of Medical Sciences. The work has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments in humans.

2.2. Genotyping

Blood samples were taken from patients and healthy individuals into ethylenediaminetetraacetic acid (EDTA) containing tubes and genomic DNA was extracted from peripheral leukocytes applying the phenol-chloroform method [12]. The extracted DNA was stored at -20°C until the proper time for the experiments. Genotyping was performed using StepOnePlus Real-Time PCR System and TaqMan MGB-based allelic discrimination method (both Applied Biosystems, Foster City, CA, USA) for rs12617656 and Amplification Refractory Mutation System-Polymerase Chain Reaction (ARMS-PCR) for rs12379034. The ARMS-PCR technique [13] is a precise and rapid method, detecting a specific SNP in a large number of SNPs and is based on a mutation or mismatched nucleotide(s) at 3' end of the primer. Three primers including two forward (mutant and wild-type) and one reverse common primer (Table 1) were designed by Primer 3 online tool (<http://primer3.ut.ee>). The ARMS-PCR was conducted in a 15 μL reaction containing 2 μL DNA template, 5U Taq DNA polymerase (SinaClon, Iran), 2.5 μL 10 \times PCR buffer (SinaClon, Iran), 0.25 μL 10 mM dNTP (SinaClon,

Table 1

Primers used in ARMS-PCR for *CDK5RAP2* gene rs12379034 genotyping, amplicon size and melting temperature of each reaction.

Primer	Sequence	Amplicon size (bp)	Temp ($^{\circ}\text{C}$)
NF	AGCTAGTCCATGAGGACAAATga	220	57.4
MF	AGCTAGTCCATGAGGACAAATgg	220	57.4
RC	CTTACTTGATCCGTCCACA	220	58.5
Seq	AGGAATCAGAAGTCCAGAA	282	58.7

NF, Normal Forward primer; MF, Mutant Forward primer; RC, Reverse Common primer; Seq, sequence primer.

Iran) and 1 μL of each specific primer (10 pmol/ μL). Two internal control primers with 0.12 μL volume in each reaction was also utilized. The following primers were used as internal control, forward: 5' TGCTCAGCCAGTTTACTTA 3' and reverse: 5' CCTGCAGG-TATATTTGGCG 3'. PCR was performed as follow: an initial denaturation at 94°C for 3 min, 40 cycles for DNA amplification consist of 30 s at 94°C , 30 s at 63.6°C , 30 s 72°C and a final extension at 72°C for 10 min. PCR products were electrophoresed in 2% agarose gel stained with DNA safe staining (SinaClon, Iran) and were observed through UV transilluminator (Fig. 1). We chose a few samples for sequencing to confirm the accuracy of the amplification (Fig. 2).

To perform TaqMan MGB-based allelic discrimination, all PCR reactions mixture contained approximately 25–75 ng of DNA, 5 μL Taq-Man Master Mix containing Taq DNA polymerase and dNTPs (Applied Biosystems, Foster City, USA), 0.25 μL Taq-Man Genotyping Assay mix containing primers and FAM or VIC labeled probes (Applied Biosystems, Foster City, USA), and distilled water for a final volume of 10 μL . Thermocyclic conditions of PCR were: initially 60°C for 30 s and then 95°C for 10 min, and subsequently 40 cycles of amplification (95°C for 15 s and 60°C for 1 min), and finally 60°C for 30 s.

2.3. Statistical analysis

All statistical analyses were performed using SPSS, version 21 (IBM Inc, Chicago, IL, USA). The generalized linear model (i.e. logistic regression) and chi-square tests were used to evaluate the association of alleles and genotypes of SNPs with disease status and clinical manifestations, respectively. Odds ratios (ORs) as the effect size with 95% confidence intervals (95% CI) were calculated in each group. The control group was evaluated for Hardy–Weinberg Equilibrium (HWE) in each locus using SHEsis online tool [14]. P values less than 0.05 were considered as statistically significant.

3. Results

Demographic and laboratory information of RA patients and healthy controls are shown in Table 2.

The genotype distribution of *DPP4* gene rs12617656 and *CDK5RAP2* gene rs12379034 SNPs in the healthy controls demonstrated no significant deviation from the HWE ($P=0.124$ and 0.256 , respectively). As shown in Table 3, the global major allele of T was regarded as the reference allele for rs12617656 SNP according to NCBI database. The C allele of rs12617656 SNP was distributed almost equally between patients and controls (33.7% vs. 32.4%). Therefore, this allele had no significantly different distribution between RA patients and healthy controls (OR = 0.95, CI: 0.74–1.23; $P=0.74$). On the other side, considering the TT genotype as the reference, genotypes of rs12617656 SNP did not represent significantly different prevalence between cases and controls. The CT genotype of this SNP was less represented in RA patients in comparison to healthy subjects (43.5% vs. 46.1%); hence the difference was not statistically significant (OR = 0.90,

Download English Version:

<https://daneshyari.com/en/article/8469437>

Download Persian Version:

<https://daneshyari.com/article/8469437>

[Daneshyari.com](https://daneshyari.com)