

Accepted Manuscript

Title: Traits allowing *Avicennia marina* propagules to overcome seawater salinity

Authors: Barhoumi Zouhaier, Ahmed Abdallah Hussain, Kamel Ahmed Saleh



PII: S0367-2530(18)30162-2
DOI: <https://doi.org/10.1016/j.flora.2018.03.003>
Reference: FLORA 51249

To appear in:

Received date: 5-11-2017
Revised date: 1-3-2018
Accepted date: 3-3-2018

Please cite this article as: Zouhaier, Barhoumi, Hussain, Ahmed, Abdallah, Saleh, Kamel Ahmed, Traits allowing *Avicennia marina* propagules to overcome seawater salinity. *Flora* <https://doi.org/10.1016/j.flora.2018.03.003>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Traits allowing *Avicennia marina* propagules to overcome seawater salinity

Barhoumi Zouhaier ^{ab}, Ahmed Abdallah Hussain^a, Kamel Ahmed Saleh^a

^a King Khalid University, Biology Department, P.O. Box- 9004, Abha- 61413 Kingdom of Saudi Arabia.

^b Laboratory of Extremophile Plants, Biotechnology Center of Borj Cedria, University Tunis El Manar, B.P. 901, Hammam-lif, Tunis, Tunisia.

*** Corresponding author: Zouhaier Barhoumi**

E-mail address: barhoumi_zou2000@yahoo.fr

Phone number: +216 79 325 848

Fax number: +216 79 325 638

Highlights

- Toxic ions (Na⁺ and Cl⁻) were mainly accumulated in the propagule coat.

Download English Version:

<https://daneshyari.com/en/article/8470150>

Download Persian Version:

<https://daneshyari.com/article/8470150>

[Daneshyari.com](https://daneshyari.com)