Accepted Manuscript

Tools and techniques

Efficient Oligo nucleotide mediated CRISPR-Cas9 Gene Editing in Aspergilli

Christina S. Nødvig, Jakob B. Hoof, Martin E. Kogle, Zofia D. Jarczynska, Jan Lehmbeck, Dorte K. Klitgaard, Uffe H. Mortensen


PII: S1087-1845(18)30004-5

DOI: https://doi.org/10.1016/j.fgb.2018.01.004

Reference: YFGBI 3107

To appear in: Fungal Genetics and Biology

Received Date: 26 September 2017 Revised Date: 5 January 2018 Accepted Date: 6 January 2018



Please cite this article as: Nødvig, C.S., Hoof, J.B., Kogle, M.E., Jarczynska, Z.D., Lehmbeck, J., Klitgaard, D.K., Mortensen, U.H., Efficient Oligo nucleotide mediated CRISPR-Cas9 Gene Editing in Aspergilli, *Fungal Genetics and Biology* (2018), doi: https://doi.org/10.1016/j.fgb.2018.01.004

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Efficient Oligo nucleotide mediated CRISPR-Cas9 Gene Editing in Aspergilli

Christina S. Nødvig^a, Jakob B. Hoof^a, Martin E. Kogle^a, Zofia D. Jarczynska^a, Jan Lehmbeck^b,

Dorte K. Klitgaard^c, & Uffe H. Mortensen^{a*}

ABSTRACT

CRISPR-Cas9 technologies are revolutionizing fungal gene editing. Here we show that survival of specific Cas9/sgRNA mediated DNA double strand breaks (DSBs) depends on the non-homologous end-joining, NHEJ, DNA repair pathway and we use this observation to develop a tool to assess protospacer efficiency in *Aspergillus nidulans*. Moreover, we show that in NHEJ deficient strains, highly efficient marker-free gene targeting can be performed. Indeed, we show that even single-stranded oligo nucleotides efficiently works as repair templates of specific Cas9/sgRNA induced DNA DSBs in *A. nidulans*, *A. niger*, and in *A. oryzae* indicating that this type of repair may be wide spread in filamentous fungi. Importantly, we demonstrate that by using single-stranded oligo nucleotides for CRISPR-Cas9 mediated gene editing it is possible to introduce specific point mutations as well gene deletions at efficiencies approaching 100%. The efficiency of the system invites for multiplexing and we have designed a vector system with the capacity of delivering Cas9 and multiple sgRNAs based on polymerase III promoters and tRNA spacers. We show that it is possible to introduce two point mutations and one gene insertion in one transformation experiment with a very high efficiency. Our system is compatible with future high-throughput gene-editing experiments.

^a Department of Biotechnology and Biomedicine, Technical University of Denmark, Søltofts Plads, Building 223, Kongens Lyngby, 2800, Denmark.

^b Department of Fungal Strain Technology, Novozymes A/S, Bagsværd, 2880, Denmark.

^c Department of Protein Diversity, Novozymes A/S, Bagsværd, 2880, Denmark.

^{*} To whom correspondence should be addressed. Tel: +45 4525 2701; Fax: +45 4588 4148; Email: um@bio.dtu.dk.

Download English Version:

https://daneshyari.com/en/article/8470413

Download Persian Version:

https://daneshyari.com/article/8470413

<u>Daneshyari.com</u>