

BMP2 and GDF5 induce neuronal differentiation through a Smad dependant pathway in a model of human midbrain dopaminergic neurons

Shane V. Hegarty, Aideen M. Sullivan, Gerard W. O'Keeffe

PII: S1044-7431(13)00066-3
DOI: doi: 10.1016/j.mcn.2013.06.006
Reference: YMCNE 2827

To appear in: *Molecular and Cellular Neuroscience*

Received date: 8 March 2013
Revised date: 24 May 2013
Accepted date: 25 June 2013

Please cite this article as: Hegarty, Shane V., Sullivan, Aideen M., O'Keeffe, Gerard W., BMP2 and GDF5 induce neuronal differentiation through a Smad dependant pathway in a model of human midbrain dopaminergic neurons, *Molecular and Cellular Neuroscience* (2013), doi: 10.1016/j.mcn.2013.06.006

BMP2 and GDF5 induce neuronal differentiation through a Smad dependant pathway in a model of human midbrain dopaminergic neurons.

Shane V. Hegarty, Aideen M. Sullivan*, Gerard W. O'Keeffe*.

Department of Anatomy and Neuroscience, Biosciences Institute, University College Cork, Cork, Ireland.

Address correspondence to

Dr. Gerard O'Keeffe
Phone (+353) 21 490 1305
Fax (+353) 21 427 3518
Email g.okeeffe@ucc.ie

or

Dr. Aideen Sullivan
Phone (+353) 21 420 5427
Fax (+353) 21 427 3518
Email a.sullivan@ucc.ie

*These authors contributed equally to this work.