

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

International Journal of Veterinary Science and Medicine

journal homepage: www.elsevier.com/locate/ijvsm

Full Length Article

Ameliorative effects of thymoquinone on titanium dioxide nanoparticles induced acute toxicity in rats

Khaled M.A. Hassanein^{a,b,*}, Yasmin O. El-Amir^{a,c}^a Department of Pathology, Faculty of Veterinary Medicine, Assiut University, Assiut 71526, Egypt^b Deanship of Scientific Research, Jazan University, Jazan, Saudi Arabia^c Department of MLT, Faculty of Applied Medical Science, Jazan University, Jazan, Saudi Arabia

ARTICLE INFO

Keywords:

Acute toxicity
Oxidative stress
Pathology
Thymoquinone
Titanium dioxide nanoparticles
Ultrastructure

ABSTRACT

Although the nanoparticles had a beneficial activity, it had also adverse effects as a result of generation of oxidative stress. The current study aimed to assess the ameliorative effect of thymoquinone (TQ) on titanium dioxide nanoparticles (TiO₂ NPs) induced acute toxicity in male rats. Forty-eight male rats were distributed into four equal groups (12 rats each). Group (1) received single oral dose of TiO₂ NPs (300 mg/kg), Group (2) received TiO₂ NPs and TQ (20 mg/kg), Group (3) received TQ and group (4) received only the vehicle and served as control group. TiO₂ NPs intoxicated group showed increased the level of lipid peroxidation product (LPO), aspartate aminotransferase (AST) and alanine aminotransferase (ALT) and decreased the level of antioxidants and testosterone. Vascular and degenerative changes in the liver and testes were observed by light microscopy as well as presence of TiO₂ NPs in the lysosomes by electron microscopy. Treatment with TQ revealed improvement of the biochemical parameters, histology and ultrastructure of the liver and testes. It was concluded that acute intoxication of rats with TiO₂ NPs induced adverse effect in the liver and testes. Administration of TQ has an ameliorative effect against oxidative stress induced by TiO₂ NPs intoxication.

1. Introduction

Titanium dioxide nanoparticles (TiO₂ NPs) are commonly used in many applications as household necessities, industrial, medicinal products as well as food colorant and white pigment [1,2]. Titanium dioxide nanoparticles induced genotoxicity, mutagenicity and cytotoxicity and potentially carcinogenicity [3]. TiO₂ NPs are used in industrial field as cosmetics in two crystalline forms anatase and rutile [4].

Nanoparticles induced cytotoxicity, apoptosis and oxidative stress to the liver cells [5,6]. TiO₂ NPs were observed in the hepatocytes and Kupffer cells in mice [5]. Jain et al. [3] reported that TiO₂ NPs induced mitochondrial swelling and disruption of the nuclear membrane in mammalian lung fibroblast cells. TiO₂ NPs might distribute from the lung tissue to other body tissue and induced systemic effects [7].

Black cummin seed (*Nigella sativa*) is used in traditional medicine in many countries as Greece, Turkey, Egypt and many other countries in Asia and Africa [8]. Thymoquinone (TQ) is the main component of the

Nigella sativa [9]. TQ has a powerful anti-inflammatory and antioxidant effects [10,11]. Moreover, TQ has anticancer activities through targeting the carcinogenic signaling molecules and immunomodulatory activities [12].

In the present study we aimed to estimate the oxidative stress, histopathology and ultrastructure of the liver and testis following oral administration of TiO₂ NPs to male rats and to assess the ameliorative effect of thymoquinone.

2. Materials and methods

2.1. Experimental design

Forty-eight male Sprague-Dawley (SD) rats of 200–230 g weight obtained from the animal house of Faculty of Medicine, Assiut University. The experiment was ethically approved by the committee of research ethics, Jazan University. Rats were acclimatized for 2 wks then

Peer review under responsibility of Faculty of Veterinary Medicine, Cairo University.

* Corresponding author at: Department of Pathology & Clinical Pathology, Faculty of Veterinary Medicine, Assiut University, Assiut 71526, Egypt.

E-mail address: khaledhassanein@aun.edu.eg (K.M.A. Hassanein).<https://doi.org/10.1016/j.ijvsm.2018.02.002>

Received 24 December 2017; Received in revised form 2 February 2018; Accepted 9 February 2018

2314-4599/© 2018 Faculty of Veterinary Medicine, Cairo University. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1

Biochemical analysis of oxidative stress, liver enzymes and testosterone in the serum of rat of all groups.

Parameters	Group 1	Group 2	Group 3	Group 4
LPO	9.76 ± 0.14 ^a	6.23 ± 0.17 ^b	4.73 ± 0.37 ^b	4.90 ± 0.05 ^b
TBARS	0.38 ± 0.08 ^b	0.66 ± 0.03 ^a	0.91 ± 0.05 ^a	0.90 ± 0.01 ^a
AST	40.10 ± 0.4 ^a	27.10 ± 0.9 ^b	20.56 ± 0.41 ^b	20.36 ± 0.2 ^b
ALT	82.60 ± 2.3 ^a	40.8 ± 1.7 ^b	34.10 ± 1.06 ^b	32.46 ± 0.4 ^b
Testosterone	1.35 ± 0.06 ^b	2.60 ± 0.2 ^a	3.77 ± 0.1 ^a	3.60 ± 0.2 ^a

The values are presented as mean ± SE, different superscripts in the same row were significantly different at ($P \leq 0.001$).

Table 2

The histopathological lesions observed by light microscopy and electron microscopy in the studied groups.

Lesions	Group 1	Group 2	Group 3	Group 4
<i>Liver lesions observed by Light microscope</i>				
Hemorrhage	++	-	-	-
Congestion	++++	+	-	-
Vacuolar degeneration	++++	+	-	-
Mononuclear infiltration	+++	-	-	-
Kupffer cell proliferation	+++	-	-	-
<i>Liver lesions observed by EM</i>				
Mitochondrial swelling	##	#	-	-
Fatty globules	###	#	-	-
Nanoparticles in Kupffer cells and hepatocytes	##	-	-	-
<i>Testicular lesions observed by Light microscope</i>				
Interstitial edema	++++	+	-	-
Congestion	++++	-	-	-
Degenerative changes	++++	-	-	-
<i>Testicular lesions observed by EM</i>				
Presence of nanoparticles	###	-	-	-

Presence of lesions by light microscopy expressed by; - (0 rats), + (1-3 rats), ++ (4-6 rats), +++ (7-9 rats), ++++ (10-12 rats). Presence of lesions by EM expressed by; - (0 rats), # (1 rats), ## (2 rats), ### (3 rats).

divided into 4 even groups (12 rats each). Group (1) received single oral dose of TiO₂ NPs (300 mg/kg b.w.) in 1% Tween 80 by stomach tube, Group (2) received TiO₂ NPs and TQ daily oral dose (20 mg/kg b.w. in corn oil by stomach tube), Group (3) received TQ daily oral dose and group (4) received single dose of the vehicle (1% Tween 80) by stomach tube. All chemical in this study were purchased from Sigma-Aldrich, Germany.

2.2. Biochemical analysis

Serum samples were collected for estimation of lipid peroxidation (LPO), total antioxidants, alanine aminotransferase (ALT), aspartate aminotransferase (AST) and testosterone levels by using commercial kits.

2.3. Histopathological analysis

Tissue samples from the liver and testis of male rats were collected after 14 days. The samples were fixed in 10% buffered formalin then routine processing, stained by hematoxylin and eosin (HE) and examined by light microscope [13].

2.4. Electron microscopy analysis

Samples from the liver and testis were obtained and immersed in

glutaraldehyde solution for transmission electron microscopy (TEM). Specimens were then processed and ultrathin sections stained with uranyl acetate and lead citrate were examined by TEM (JEOL100 CXII, Japan) in the EM Unit, King Khalid University, KSA [13].

2.5. Statistical analysis

All values were presented as mean ± SE for the studied parameters and analyzed using GraphPad Prism software program.

3. Results

3.1. Biochemical results

The levels of lipid peroxidation (LPO) and liver enzyme markers (AST & ALT) were significantly increased ($P \leq 0.001$) in TiO₂ NPs intoxicated group compared to other groups. Conversely, testosterone and total antioxidants levels were decreased significantly. Administration of TQ recovered the serum levels of LPO, total antioxidants, AST, ALT and testosterone (Table 1).

3.2. Histopathology

The lesion score of the histopathology of the studied groups were summarized in Table 2. Histopathological examination of the liver tissue of the rat intoxicated with TiO₂ NPs revealed pathological alterations. Congestion of the central veins and dilatation of the hepatic sinusoids were observed (Fig. 1a). Focal hemorrhagic areas were observed in the hepatic parenchyma as well as coagulative necrotic changes in the hepatocytes. Proliferation of Kupffer cells were also seen (Fig. 1b). Peri-central aggregations of mononuclear cells especially lymphocytes were consistent phenomena in the examined liver in group 1 (Fig. 1c). Vacuolar degeneration of some hepatocytes was observed (Fig. 1d). Some examined sections revealed TiO₂ NPs in Kupffer cells which appeared as eosinophilic particles (Fig. 1e). Examined liver section from TQ treated group showed normal architecture with dilatation of the central veins (Fig. 1f). The liver of control group showed normal histological appearance (Fig. 1g).

Examination of testis in rats intoxicated with TiO₂ NPs revealed congestion of the blood vessels (Fig. 2a). Interstitial edema was observed and the seminiferous tubules appeared atrophied. The seminiferous tubules showed degenerative changes in which Sertoli cells appeared necrosed with few germinal cells (Figs. 2b, c). TQ reduced the histopathological alteration induced by TiO₂ NPs (Fig. 2d). No histopathological changes could be seen in control group (Fig. 2e).

3.3. Transmission electron microscopy

The lesion score of EM examination in the studied groups were summarized in Table 2. With ultrathin sections, the cytoplasm of hepatocytes intoxicated with TiO₂ NPs revealed mitochondrial swelling (Fig. 3a). Numerous intracytoplasmic fatty globules of variable sizes were observed associated with pyknotic mitochondria with intense metrical condensation (Fig. 3b). Kupffer cells revealed several lysosomes loaded with TiO₂ NPs which appeared electron dense granules either minute or compact particles. These particles were also seen in the cytoplasm of hepatocytes (Fig. 3c). Examined hepatocytes from TQ treated group showed normal nucleus and cytoplasmic organelles as well as few intracytoplasmic fatty globules (Fig. 3d). The hepatocytes from control group showed normal nucleus and cytoplasmic organelles (Fig. 3e). Examination of the ultrathin section of seminiferous tubules in TiO₂ NPs group showed abundant lysosomes loaded with TiO₂ NPs

Download English Version:

<https://daneshyari.com/en/article/8482167>

Download Persian Version:

<https://daneshyari.com/article/8482167>

[Daneshyari.com](https://daneshyari.com)