



Full Length Article

Serum cortisol of Sahel goats following rumenotomy with assorted anaesthetics and sutures



Abubakar Mshelia Saidu^{a,*}, Paul B. Bokko^a, Abdullahi Mohammed^a,
David N. Bukbuk^b, Ephraim Igwenagu^c

^a Department of Veterinary Surgery and Radiology, University of Maiduguri, Maiduguri, Borno State, Nigeria

^b Department of Microbiology, University of Maiduguri, Maiduguri, Borno State, Nigeria

^c Department of Veterinary Pathology, University of Maiduguri, Maiduguri, Borno State, Nigeria

Received 17 March 2016; revised 14 October 2016; accepted 16 October 2016

Available online 30 November 2016

KEYWORDS

Sahel goat;
Cortisol;
Rumenotomy;
Polyglycolic acid;
Chromic catgut;
Sutures;
Stress

Abstract The utmost need for pragmatic combination of surgical sutures and local anaesthetic that would evoke minimal post-surgical stress response and allow uncomplicated healing is essential for successful surgeries. Fifteen Sahel goats were randomly allocated into three groups A, B and C to quantitatively assay (ELISA) serum cortisol profiles following rumenotomy, as markers of surgical stress. Diazepam at 0.2 mg/kg was administered intravenously to groups A and B with subsequent lidocaine HCl and bupivacaine inverted-L block respectively. Group C did not receive any treatment. Chromic catgut (CCG) and polyglycolic acid (PGA) sutures were used for rumen and abdominal muscles closure for groups A and B respectively and nylon for skin closure. Blood samples were taken at post anaesthetic induction (PAI) and post-surgery at 0, 5, 8, 24, 48 and 72 h. The Group A goats expressed serum cortisol that was significantly high 52.76 ± 6.12 ng/mL at 5 h post-surgery. At 8 h post-surgery serum cortisol for both groups A (72.53 ± 3.79 ng/mL) and B (61.59 ± 3.90 ng/mL) were at their peak. Serum cortisol levels compared to the baseline data were significantly different ($P < 0.05$) at 5, 24, and 48 h for the CCG goats. The serum cortisol levels at 72 h drastically decreased to 20.53 ± 8.74 ng/mL for groups A and 17.59 ± 2.45 ng/mL for group B and were not significantly different ($p > 0.05$). Cortisol responses unambiguously indicate that diazepam-bupivacaine induce less stress than Diazepam-lidocaine, hence a preferred anesthesia. Moreover, polyglycolic acid sutures are associated with less inflammatory reaction than chromic catgut.

© 2016 Faculty of Veterinary Medicine, Cairo University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Cortisol is an important steroid hormone secreted by the adrenal cortex and associated with inflammation, carbohydrate

metabolism and stress reaction [1]. Stress is a disruptive event accompanied by predictable biochemical, physiological, cognitive and behavioral changes that are directed either toward altering the stressful event or accommodating to its effects [2]. It refers to external and internal forces (stressors) applied to animals or other biological systems, leading to measurable alteration of a physiological steady state. These changes termed stress response comprise of metabolic, immunological and neuroendocrine changes following injury or trauma [3,4].

* Corresponding author.

E-mail address: abubakarsm51@gmail.com (A.M. Saidu).

Peer review under responsibility of Faculty of Veterinary Medicine, Cairo University.

<http://dx.doi.org/10.1016/j.ijvsm.2016.10.008>

2314-4599 © 2016 Faculty of Veterinary Medicine, Cairo University. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Stressors are classified as physical/environmental or psychologic/perceived [5]. These stressors influence activation of the hypothalamic pituitary adrenal axis followed by synthesis of corticotrophin or adrenocorticotrophic hormone (ACTH) by the anterior pituitary. This stimulates adrenal cortical secretion of glucocorticoids leading to increased circulating concentration of cortisol. Surgery is one of the most potent activators of this process among others such as infection, tissue injury, trauma, neoplastic growth or immunological disorder [3,5]. Psychological and physical factors like anxiety and pain influence the activation of the hypothalamic–pituitary–adrenal axis (HPA) leading to cortisol release; a core mediator of stress response [3,6] which in turn promotes glycogenolysis and gluconeogenesis resulting in hyperglycemia. The stress induced cortisol secretion exists so that the body can handle and recuperate after an injury, physical activity or physiological strain [3]. Animals respond differently to various kinds of stressors. Plasma cortisol is a useful indicator of stress and muscle damage and data are very limited on responses to surgical stress in veterinary medicine [7]. Studies have shown elevated cortisol concentrations above normal 18 ng/mL levels due to isolation of goats from their social group as emotional stress [7]. When social isolation was not combined with holding treatment, the stress levels were significantly lower in goats that were able to maintain visual contact with other animals [7]. Stressors such as transportation significantly increases plasma cortisol levels within 30 min after beginning of transportation and reached a peak value at one hour in adult male goats; in addition, young goats had higher plasma cortisol concentrations than older ones [7]. The stress indicators and markers elucidate the prospective direction of stressors. These aid in selection of anaesthetic agents, sutures and other corrective strategies that will elicit minimum stress and optimum reparative pathways. For instance, local/regional anaesthesia for rumenotomy that helps in reduction of pain and desirable pathophysiological responses [3,8,9], as well as the choice of suture materials that greatly dictates post-surgical tissue reactions and inflammatory processes. Therefore, “there is a practical need for pragmatic combination of surgical sutures and local anaesthetic that would evoke minimal post-surgical stress response and allow uncomplicated healing”. Sutures and anaesthetic agents putatively produce distinctive tissue reaction and inflammatory process. However, the indicators and biomarkers of surgical stress patterning following surgery in Sahel goats remain unclear. This study aims to evaluate the stress induced by induction of local anesthesia and rumenotomy in the Sahel goat by determining the levels of serum cortisol in Diazepam-Lidocaine Combination (DLC) and Diazepam-Bupivacaine Combinations in the event of choice of chromic catgut (CCG) and polyglycolic acid (PGA) sutures.

2. Materials and methods

2.1. Animals

Fifteen apparently healthy Sahel goats aged 15.10 ± 4.12 months weighing 17.70 ± 3.3 kg bw were procured from Maiduguri livestock market in Borno state, Nigeria. Maiduguri is the Capital of Borno state in the northeastern region of Nigeria. Maiduguri is located within latitude 11.15°N and

longitude 30.05°E the Sudano-Savanna zone [10]. Clearance from the Faculty of Veterinary Medicine Animal Ethics Committee, University of Maiduguri, Nigeria was obtained for the research. Using a random number generator (RNG®), the goats were randomly allocated to 3 groups of 5 goats A, B and C, with A and B as experimental groups while C as a control group.

2.2. Preoperative preparation

The goats were fasted for 12 and 6 h for feed and water respectively. The left paralumbar fossa was clipped and the area was aseptically prepared with 0.2% chlorhexidine gluconate (Savlon®, Vervaaingdeur, Johnson and Johnson (pty) Ltd, London) (60 mL stock per liter of sterile water). Prior to anaesthesia and surgery, a drip infusion line was instituted via the jugular vein and commenced fluid therapy with 0.9% normal saline solution (Juhel®, Fabrique par Juhel Nig. Ltd/Awka, Anambra, Nigeria) as maintenance fluid, set at 180 mL per hour (1 drop per second). The experimental goats were sedated with Diazepam, (SJG®, FazulEllahie, Pvt Ltd, Karachi, Pakistan), given at a dose rate of 0.2 mg/kg IV. The goats in group A received 2% Lidocaine HCl 4 mg/kg (Lidocaine®, Kwaliti Pharmaceuticals (P) Ltd. NagKalan, Majitha Road, Amritsar, INDIA) to create an inverted L-block regional anaesthesia on the left flank (Paralumbar fossa) immediately after sedation. Animals in group B received 0.5% Bupivacaine 1.5 mg/kg (Marcaine®, Astrazeneca PLC, IngiItire Lisansiile, Istanbul, Turkey), for inverted L-block post Diazepam, (SJG®, Fazul Ellahie, Pvt Ltd, Karachi, Pakistan), sedation. Five minutes after induction, 5 mL of blood samples were drawn from the opposite jugular vein into a plain vacutainer tubes and kept at room temperature for 2 h before centrifugation at 4000×g for 3 min in an electronic centrifuge (Centrifuge 800B®, Union Laboratories, England).

2.3. Surgery

The standard laparotomy method ‘through-and-through’ incision was made on the upper paralumbar fossa of the left flank [11]. The rumen was gently pulled out of the incision and firmly anchored to the skin ventrally and dorsally with two of six towel clamps for the technique. A ten centimeter (10 cm) incision was made over less vascular portion of the rumen greater curvature and its edges were fixed caudally and cranially to the skin incision with towel clamps [12]. The remaining clamps were used to secure the rumen edges between the previously placed clamps with their handles away from the incision site. The rumen edges covered the skin edges by 2–3 cm. The rumen were explored and foreign materials mostly plastic bags weighing 1.26 ± 0.18 and 1.44 ± 0.86 kg for groups A and B respectively were removed, 0.9% saline solution was used to rinse the rumen ingesta. To commence rumen closure, the caudal and cranial clamps were removed first leaving the dorsal and ventral clamps in place. Double layer Cushing suture pattern was used to invert the rumen edges with a number 2 chromic catgut (LIFECARE®, Anhui Kangning Industrial group Co. Ltd, Tianchang City, Anhui, China) and polyglycolic acid sutures (Atramat®, International Farmaceutica, Planta, Mexico), for groups A and B respectively.

Download English Version:

<https://daneshyari.com/en/article/8482434>

Download Persian Version:

<https://daneshyari.com/article/8482434>

[Daneshyari.com](https://daneshyari.com)