

## Disseminated cryptococcosis in a HIV-negative patient: Case report of a newly diagnosed hypertensive adult presenting with hemiparesis

Raymond M. Wilson<sup>a,1</sup>, Nyambura Moremi<sup>b,1</sup>, Martha F. Mushi<sup>b,\*</sup>, Oliver Bader<sup>c</sup>, Patrick S. Ngoya<sup>d</sup>, Bernard M. Desderius<sup>a</sup>, Peter Rambau<sup>e</sup>, Rodrick Kabangila<sup>a,f</sup>, Uwe Groß<sup>c</sup>, Stephen E. Mshana<sup>b</sup>

<sup>a</sup> Department of Internal Medicine, Bugando Medical Centre/Catholic University of Health and Allied Science, Mwanza, Tanzania

<sup>b</sup> Department of Microbiology and Immunology, Weill Bugando School of Medicine, Catholic University of Health and Allied Sciences, Mwanza, Tanzania

<sup>c</sup> Institute of Medical Microbiology, University Medical Center Goettingen, Germany

<sup>d</sup> Department of Radiology, Bugando Medical Centre, Mwanza, Tanzania

<sup>e</sup> Department of Pathology, Weill Bugando School of Medicine, Mwanza, Tanzania

<sup>f</sup> Kamanga Medics Hospital, Mwanza, Tanzania

### ARTICLE INFO

#### Keywords:

Disseminated cryptococcosis

*C. gattii*

Meningism

*Cryptococcus deuterogattii*

HIV negative

### ABSTRACT

We report a case of disseminated cryptococcosis in a 42-year old immunocompetent female. Prior to admission at Bugando Medical Center, the patient was attended at three hospitals for hypertension and clinically diagnosed malaria. Following diagnosis of disseminated *Cryptococcus* at our center, she was successfully treated with fluconazole but remained with visual loss. Blood cultures should be considered in the management of any adult presenting with fever to enable early detection of the least expected differentials like in this case.

### 1. Introduction

Cryptococcal meningitis (CM) is among the common AIDS-defining illnesses, and the leading cause of AIDS-related death among HIV-infected patients admitted at Bugando Medical Center (BMC) with mortality of up to 67% [1]. In contrast to *C. neoformans* which is prevalent worldwide among HIV-infected patients [2], *C. gattii* was recently identified as a species found commonly in tropical regions infecting immunocompetent people presenting with neurological and pulmonary cryptococcoma [3]. Cases of disseminated cryptococcosis in immunocompetent individuals have rarely been reported in Sub-Saharan Africa partly due to limited diagnostic facilities and low index of suspicion of the attending clinicians, and this could be a result of overlap of signs and symptoms with other diseases [4,5]. We present a disseminated cryptococcosis in a HIV-negative woman, a newly diagnosed patient with hypertension who presented with right-sided hemiparesis, fever and signs of meningism. This case stresses the importance of high index of suspicion and extensive work-up in immunocompetent patients presenting with meningism and fever in a resource-limited setting.

### 2. Case

In April 2017, a 42-year-old woman, presented to the Bugando Medical Center (BMC) as a self-referred patient from home (Mahina, Mwanza) following persistence of headache for one month, fever for two weeks and right-sided weakness for one week. The headache started gradually and progressively became more severe with time. The headache was more severe on the frontal region, throbbing in nature, associated with blurred vision and persistent fevers. She reported to have episodes of nausea and vomiting. The right-sided body weakness started gradually one week prior to BMC admission but the patient could still walk with minimal support. On the course of these symptoms, the patient received care at three hospitals in Mwanza city where she was diagnosed to have hypertension and symptomatic malaria and kept on losartan H. 62.5 mg daily, antimalarials and antibiotics of which she could not recall the name. She denied any history of cough, convulsions, loss of consciousness, head trauma, diabetes mellitus, and steroid/chemotherapy use. On examination, she had fever of 38.1 °C, blood pressure of 180/100 mmHg, positive Kerning's and Brudzinski's signs, was photophobic with pupils that were consensually reacting to light, decreased visual acuity bilaterally, and showed decreased deep reflexes with power grade of 4/5 on right arm and leg. The patient was

\* Corresponding author.

<sup>1</sup> Contributed equally.



Fig. 1. Firm, non-tender and non-pulsating nodule on the right nostril.

also found to have a firm, non-tender and non-pulsating nodule ( $9 \times 6$  mm) on the right nostril (Fig. 1). Respiratory, abdominal and fundoscopic examination were uneventful.

Based on examination findings, the patient was clinically diagnosed to have an ischemic stroke with right-sided hemiparesis due to hypertension with a differential diagnosis of bacterial meningitis. She was started on high dose ceftriaxone i.e. 2 g i.v. 12 hourly and ampicillin 2 g i.v. 6 hourly as the empirical management of bacterial meningitis.

A complete blood count revealed white blood cell count, granulocytes, lymphocytes, platelet count and hemoglobin to be  $7.0 \times 10^9$ /U/L, 70%, 23.8%,  $373 \times 10^9$  U/L and 10 g/dL, respectively. HIV testing according to Tanzania HIV Rapid Test Algorithm of 2015 was negative. The CD4 and CD8 cell counts were 519 and 299 cells/ $\mu$ L, respectively. Renal and liver function tests were all within a normal range. Moreover, a chest X-ray was also normal.

On day 3, CT imaging was requested to rule out space occupying lesions, and the findings showed multifocal hypodense intracerebral lesions (not shown) thought to be subacute hemorrhagic infarcts secondary to an underlying vasculitis or a thromboembolic phenomenon, hence a lumbar puncture was done. The CSF had 35 cells/mm<sup>3</sup>, positive *Cryptococcus* antigen test, and positive Indian and Pandy's test. As a way of ruling out autoimmune vasculitis, Perinuclear Anti-Neutrophil Cytoplasmic Antibodies (p-ANCA), Cytoplasmic-ANCA, and Antineutrophil Cytoplasmic Antibodies (ANA) profiles were performed which were all negative. Carotid Doppler ultrasound was also done for the same purpose and was unremarkable.

The biopsy was taken from the nostril, fixed in 10% formalin and

sent for histological examination. The sample was processed and embedded in paraffin and section was stained by hematoxylin and eosin. Histological findings of the biopsied nodule revealed a skin with underlying chronic inflammatory cell infiltrate with foamy macrophages (Fig. 2A), scattered round fungal bodies with a thick capsule and a halo indicative of fungal organisms (Fig. 2B). Neither granuloma nor giant cells could be identified. The fungal bodies were positive for Periodic Acid Schiff (not shown). The diagnosis of fungal infection suggestive for *Cryptococcus* spp. was made.

Until day 5 the patient was still on the management of bacterial meningitis with no improvement. The CSF culture, India ink stain and *Cryptococcus* antigen test on this day confirmed the cryptococcal infection and the patient was started on CM management as per BMC protocol with 1.2 g i.v. fluconazole (induction), 8 weeks of 800 mg p.o. fluconazole (consolidation), and a maintenance dose of 400 mg p.o. for one year. As part of CM management protocol, serial lumbar punctures were performed on day 0 of CM diagnosis, day 1, 3 and 7 and 14 to control the management and reduce intracranial pressure (ICP) of which at the 14th day post i.v. fluconazole, the CSF culture remained negative.

Antifungal susceptibility was tested using broth microdilution method according to EUCAST guidelines [6]. The isolate was susceptible to fluconazole (0.063  $\mu$ g/ml), caspofungin (0.004  $\mu$ g/ml), micafungin (0.004  $\mu$ g/ml), posaconazole (0.008  $\mu$ g/ml) and voriconazole (0.031  $\mu$ g/ml). The molecular characterization using URA3-RFLP typed the isolate as *Cryptococcus deuterogattii* (genotype AFLP6/VGII) among the newly categorized species of *C. gattii* [7]. The source of the agent remained unclear, as the patient did not recall any exposure to bird droppings, or to eucalyptus (or similar) trees.

After 14 days of induction with i.v. fluconazole, the patient developed seizures upon which MRI was requested and demonstrated multiple circumscribed T2 and FLAIR hyperintense signal, FLAIR hyperintense with internal nodular and peripheral capsule isointense signal, T1 hypointense cystic lesions with TI post-contrast nodular enhancement, suggestive of infective abscesses or granulomas (Fig. 3A).

Five months later, the patient was seen at the clinic; she was on her maintenance dose with only a single complaint of blurred vision of which one year later had progressed to bilateral loss of vision. One year later, a follow up MRI demonstrated almost complete resolution of most of the lesions post treatment (Fig. 3B).

### 3. Discussion

The understanding of existence of non-HIV associated CM among clinicians in Sub-Saharan Africa including Tanzania is rare due to a limited documentation of cases, as seen in this patient. The later context results into low index of suspicion [5] which in turn results into either a delay as demonstrated at a tertiary healthcare facility (BMC). In all




Fig. 2. Histological findings (A) Foamy macrophages in a background (Arrow) (B) Scattered round bodies with thick capsule and a halo (arrows) indicative of fungal cells.

Download English Version:

<https://daneshyari.com/en/article/8484599>

Download Persian Version:

<https://daneshyari.com/article/8484599>

[Daneshyari.com](https://daneshyari.com)