

Subcutaneous abscesses caused by *Trichophyton rubrum* in the unilateral groin of an immunocompromised patient: A case report

Utako Okata-Karigane^{a,*}, Yasuki Hata^b, Emiko Watanabe-Okada^a, Shunichi Miyakawa^a, Michi Ota^c, Yutaka Uzawa^d, Shigekazu Iguchi^d, Atsushi Yoshida^d, Ken Kikuchi^d

^a Department of Dermatology, Kawasaki Municipal Hospital, 12-1 Shinkawadori, Kawasaki, Kanagawa 210-0013, Japan

^b Department of Dermatology, Saiseikai Yokohamashi Tobu Hospital, 3-6-1 Shimosueyoshi, Tsurumi, Yokohama, Kanagawa 230-8765, Japan

^c Department of Dermatology, Showa University Hospital, 1-5-8 Hatanodai, Shinagawa, Tokyo 142-8666, Japan

^d Department of Infectious Diseases, Tokyo Women's Medical University, 8-1 Kawada-cho, Shinjuku, Tokyo 162-8666, Japan

ARTICLE INFO

Keywords:

Dermatophyte infection
Deeper dermal dermatophytosis
Abscess
Granuloma
Trichophyton rubrum

ABSTRACT

A 60-year-old Japanese man presented with multiple subcutaneous nodules in his left groin. Histologically, the nodules consisted of suppurative granulomas and abscesses not involving the hair follicles. *Trichophyton rubrum* TWCC57922 was detected by fungal culture and polymerase chain reaction (PCR) sequencing of the rDNA genes. We diagnosed these nodules as deeper dermal dermatophytosis, a rare form of invasive dermatophytosis. He was treated with terbinafine. We compared these findings with previous reports of deep dermal dermatophytosis.

1. Introduction

Dermatophytes usually infect the keratinized layers of the epidermis, hair follicles, and nails. In rare cases, dermatophytes proliferate in the deep dermis and subcutaneous tissues. Although there is no specific classification for invasive dermatophytosis, it could be divided into three forms: Majocchi's granuloma, deeper dermal dermatophytosis, and disseminated dermatophytosis [1]. Deeper dermal dermatophytosis is rare and presents with subcutaneous granulomas or abscesses, with or without the involvement of hair follicles. We describe the case of an immunocompromised patient with multiple subcutaneous abscesses in the left groin caused by *Trichophyton rubrum*, and treated with terbinafine.

2. Case

A 60-year-old Japanese man developed a subcutaneous nodule in his left groin on day – 169 and visited our department on day – 147. We performed an incision and drainage of the nodule, which evacuated yellowish-colored, purulent material. Bacterial cultures on blood agar (POURMEDIA; horse, Eiken Chemical, Tokyo, Japan) and Bromthymol Blue (BTB) lactose agar (POURMEDIA Modified DRIGALSKI Agar; Eiken Chemical, Tokyo, Japan), and fungal cultures on M Green Yeast and Mold medium (MGYM) agar (Kyokuto, Tokyo, Japan) were all negative. The wound healed in 1 month. Around day – 90, the patient noticed

that the nodule had returned. It gradually enlarged, and other nodules began to appear around the same site. He visited our department again on day 0.

He had been receiving 10 mg of prednisolone and 4 mg of tacrolimus for interstitial pneumonia associated with polymyositis for 10 years. He also had a history of psoriasis vulgaris.

The physical examination on day 0 revealed multiple well-demarcated, elastic, soft, mildly tender, subcutaneous nodules in the left groin measuring 10–20 mm (Fig. 1a). One of the nodules was fluctuant and subsequently evacuated yellow purulent material. No bacterial or fungal cultures were obtained at that time.

Laboratory tests revealed a white blood cell count of 12,170/μl (normal range, 3900–9800/μl), a C-reactive protein level of 0.79 mg/dl (< 0.30 mg/dl), and a beta-D-glucan level of 272 pg/ml (< 20.0 pg/ml). Multiple subcutaneous nodules were identified on a computed tomography (CT) scan (Fig. 1b).

We performed a partial excision of the nodules on day + 82. The histopathological examination showed a well-demarcated nodule comprising multiple abscesses and suppurative granulomas with fibrous septae (Fig. 2a). The suppurative granulomas consisted of a large number of neutrophils surrounded by histiocytes and giant cells (Fig. 2b). There were many spore chains and septate hyphae in the abscesses and granulomas, which were more obvious in the Grocott-stained section (Fig. 2c).

We changed the MGYM agar to BD BBL™ Sabouraud Dextrose Agar

* Corresponding author.

E-mail address: utakokarigane@gmail.com (U. Okata-Karigane).

<https://doi.org/10.1016/j.mmcr.2018.03.001>

Received 18 December 2017; Received in revised form 10 February 2018; Accepted 1 March 2018

Available online 06 March 2018

2211-7539/© 2018 The Authors. Published by Elsevier B.V. on behalf of International Society for Human and Animal Mycology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. Clinical and radiographic pictures of skin lesions. (a) Multiple elastic, soft, subcutaneous nodules in the left groin. (b) Multiple subcutaneous nodules on a CT scan.

with Chloramphenicol and Gentamicin (Becton, Dickinson and Company, Japan), and the culture finally yielded a white, downy, yellowish reverse colony (Fig. 3a). The fungal organism was identified as *T. rubrum* TWCC57922 by polymerase chain reaction (PCR) sequencing of the ribosomal region spanning the internal transcribed spacers (ITS1 and ITS2) and the 5.8S rRNA following the protocol described by Hir-otsu et al. (Fig. 3b) [2]. The 604 bp of ITS1–5.8S rDNA–ITS2 region of TWCC 57992 was completely identical (100%) to that of each *T. rubrum* strain including the type strain CBS 392.58.

After the partial excision on day + 82, the nodules, which had been diminishing, began to gradually regrow around day + 300. The patient was started on a regimen of oral terbinafine 250 mg daily on day + 355, and the nodules were almost resolved after 3 months of therapy. He continued his treatment for a total of 6 months and did not be detected any nodules by a CT scan. There were no side effects during the treatment period.

3. Discussion

Dermatophytosis usually affects the stratum corneum, the hair follicles, and the nails. Deeper dermatophytosis that invades the deep dermis or subcutis is rarely seen. Although there are no specific classifications for invasive dermatophytosis, Marconi et al. have classified this condition into three forms: Majocchi's granuloma, deeper dermal




Fig. 2. Histopathology of the surgical specimen. (a) The nodule composed of abscesses and granulomas with fibrous septae. (Hematoxylin-eosin stain; loupe) (b) The suppurative granuloma consisted of neutrophils surrounded by histiocytes and giant cells. (H&E; original magnification: X100.) (c) Spore chains and septate hyphae in the granuloma. (Grocott stain; original magnification: X100.).

dermatophytosis, and disseminated dermatophytosis [1]. Majocchi's granuloma is the most common form, in which dermatophytes proliferate in the dermis through the hair follicles to form granulomas [3]. Deeper dermal dermatophytosis is rarer and is detected as an abscess, a suppurative granuloma, or a pseudocyst, with or without the involvement of hair follicles. Disseminated dermatophytosis can involve various skin sites and internal organs.

We diagnosed this patient as having deeper dermal dermatophytosis because he presented with subcutaneous abscesses and suppurative

Download English Version:

<https://daneshyari.com/en/article/8484675>

Download Persian Version:

<https://daneshyari.com/article/8484675>

[Daneshyari.com](https://daneshyari.com)