

The important role of co-infections in patients with AIDS and progressive disseminated histoplasmosis (PDH): A cohort from Colombia

Diego H. Caceres^{a,*}, Ángela M. Tobón^{a,b}, Ángela Restrepo^a, Tom Chiller^c, Beatriz L. Gómez^{a,d}

^a Medical and Experimental Mycology Unit, Corporación para Investigaciones Biológicas (CIB), Medellín, Colombia

^b Hospital La Maria, Medellín, Colombia

^c Mycotic Diseases Branch, Centers for Disease Control and Prevention, (CDC), Atlanta, United States

^d School of Medicine and Health Sciences, Universidad del Rosario, Bogotá, Colombia

ARTICLE INFO

Keywords:

AIDS
Histoplasmosis
Histoplasma
Co-infection

ABSTRACT

A total of 23/45 (51%) patients with AIDS and histoplasmosis from Medellín, Colombia had other infections. Tuberculosis was the most common ($n = 16/23$, 70%). Pneumocystosis and cryptococcosis were found in three patients each (13%), bacterial infection and cytomegalovirus occurred each in two patients (9%) while toxoplasmosis, herpes virus and esophageal candidiasis were recorded in one patient each (4%). Of all co-infected patients, 18/23 (78%) had one, four (17%) had two and one (4%) had three additional opportunistic infections.

1. Introduction

Histoplasmosis is a common infection in the Americas, although it is diagnosed in certain countries of Asia (China, India, Malaysia, Indonesia, Singapore, Thailand, Vietnam and Japan) and Africa (Democratic Republic of Congo, Congo, Liberia, Ivory Coast, Gambia, Uganda, Central African Republic, Guinea Bissau and Senegal) [1–3]. The clinical presentation depends partly on the concentration of infectious particles in the inoculum, as well as on the condition of the host's immune system at the time of infection [4,5]. Highly susceptible populations are recognized and include individuals at the extremes of life, persons undergoing immunosuppressive therapy, solid organ recipients and, especially, patients infected with the human immunodeficiency virus (HIV) who have CD4 T cells counts below 150 per μL [4].

In persons living with HIV/AIDS (PLWHA), the mycosis often produces a severe clinical form of disease called progressive disseminated histoplasmosis (PDH), which has a high mortality if treatment is not initiated promptly [4,5]. As has been previously reported, PDH symptoms are nonspecific, and among PLWHA, symptoms may be similar to those observed in other infectious diseases, thus complicating diagnosis and treatment [4,6–8]. In Colombia and other endemic countries in the Americas, where access to antiretroviral therapy (ART) can be limited, histoplasmosis is a cause of high morbidity and mortality (up to 30%) in PLWHA [9].

2. Cases

An observational, cross-sectional study of patients with histoplasmosis was performed between May 2008 and August 2011 in a cohort of PLWHA from La Maria Hospital in Medellín, Colombia [8]. The diagnosis of histoplasmosis was made based on the guidelines of the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the Mycoses Study Group (EORTC/MSG) [10]. A diagnosis was considered proven if *H. capsulatum* was isolated from any of the following samples: blood, tissue, sterile fluids or respiratory specimens or by a positive histopathological analysis, with presence of intracellular yeasts, identified by Wright staining. A diagnosis was considered probable if *H. capsulatum* could not be isolated, but there was evidence of reactive serology such as presence of either H or M precipitin band or both bands by the immunodiffusion (ID) test and a titer of 1:32 or higher with histoplasmin antigen by the complement fixation (CF) test, or a positive antigenuria test [10]. Co-infections present in these patients were defined based on clinical, radiological and microbiological criteria.

A standardized data collection form was developed in Microsoft Access consisting of variables for clinical findings, laboratory testing, treatment and outcome. For statistical analysis, absolute and relative frequencies were calculated and tested for normality. To identify statistical association, we used the Chi2 test, Student *t*-test or Mann-Whitney *U* test. In order to identify differences among histoplasmosis patients with and without co-infections, statistical associations were defined by a *p* value less than 0.05. All analyses were performed using

* Corresponding author.

E-mail address: diegocaceres84@gmail.com (D.H. Caceres).

<http://dx.doi.org/10.1016/j.mmcr.2017.07.004>

Received 23 November 2016; Received in revised form 25 January 2017; Accepted 18 July 2017

2211-7539/© 2017 The Author(s). Published by Elsevier B.V. on behalf of International Society for Human and Animal Mycology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. Study flow chart of analyzed subjects. General characteristics of 45 patients with AIDS and progressive disseminated histoplasmosis (PDH). In 22 (49%) patients no other opportunistic infection was reported at time of diagnosis of PDH; presence of co-infections were reported in the remaining 23 (51%) patients with AIDS and PDH.

the software EPIDAT 3.1 and STATA 8.0.

A total of 45 patients with AIDS and PDH were identified, 30 (67%) were classified as proven cases and 15 (33%) as probable cases. At the time of enrollment in the study, patients with histoplasmosis had the following clinical characteristics: a median CD4 cell concentration of 30 per μL (Range [R] = 3–540 cells per μL); and a median viral load of 151,914 copies per μL (R = 40–6,678,540 copies per μL) (Fig. 1); median body temperature was 37.4 °C (R = 36–39 °C); median body mass index was 18.04 (R = 12.3–24.8). A total of 39 (87%) patients with histoplasmosis had weight loss at enrollment and 18 (40%) had a Karnofsky score under 30. Gastrointestinal manifestations were present in 38 (84%) patients. Patients also had pulmonary manifestations (75%), neurological abnormalities (64%), lymphadenopathies (58%), and skin (49%) and oral mucosal lesions (44%). A total of 8 (18%) patients died.

Co-infections were diagnosed in 23 of the 45 patients (51%). There were 19 males (83%) and 4 females (17%) with a median age of 38 years (R = 21–55 years). The median CD4 T cell count was 25 cells per μL (R = 3–330 cells per μL) and the HIV viral load was 100,000 copies per μL (R = 266–6,678,540 copies per μL). No statistically significant differences among patients with and without co-infections were found (Fig. 1). There was also no statistically significant differences in mortality among patients with and without co-infections (Fig. 1).

Of the 23 patients presenting with histoplasmosis and other

infections, 18 (78%) had one additional opportunistic infection, 4 patients (17%) had two, and one patient (4%) had 3 additional infections (Fig. 2). Tuberculosis (TB) was found in 16 of 23 patients (70%) and was the main co-infection followed by pneumocystosis and cryptococcosis diagnosed in 3 of 23 patients each (13%). Bacterial infections (*Salmonella*) and *Cytomegalovirus* (CMV) occurred in 2 patients each (9%), while 1 patient exhibited concomitant toxoplasmosis, herpes virus and esophageal candidiasis. It is important to note that all 4 patients presenting with either bacterial disease or CMV infection died.

When we compared clinical manifestations grouped by systems among patients with and without co-infections, we observed differences in the presence of lymphadenopathy and neurological symptoms. All other variables analyzed did not show differences between the two groups (Fig. 3).

3. Discussion

Among this cohort of PLWHA who were identified as having progressive disseminated histoplasmosis, over half had an additional opportunistic infections [11]. This is higher than those reported in Panama (25%), French Guiana (37–42%), Argentina (42%) and Brazil (43%). However, tuberculosis was the most frequently reported opportunistic infection in all these countries [12–14,16,17].

It is clear from these results that in this patient population, there is a




Fig. 2. Distribution of co-infections observed in 45 patients with AIDS and progressive disseminated histoplasmosis (PDH). In 22 patients no other opportunistic infection was reported at time of diagnosis of PDH (49%). In the remaining 23 patients with co-infections (51%), 18 (78%) had 1 additional opportunistic disease, 4 patients (17%) had 2, and 1 patient (4%) had 3 different opportunistic diseases.

Download English Version:

<https://daneshyari.com/en/article/8484745>

Download Persian Version:

<https://daneshyari.com/article/8484745>

[Daneshyari.com](https://daneshyari.com)