

The treatment of a pregnant HIV positive patient with cryptococcal meningitis in Malawi. Case report and review of treatment options

Philip D. Bright^{a,*}, Duncan Lupiya^b, Joep J. van Oosterhout^{b,c}, Amy Chen^d, Thomas S. Harrison^a, Adrienne K. Chan^{b,e}

^a Institute of Infection and Immunity, St Georges University of London, Cranmer Terrace, London SW17 0RE, United Kingdom

^b Dignitas International, Zomba, Malawi

^c Department of Medicine, University of Malawi College of Medicine, Blantyre, Malawi

^d Trillium Health Partners, University of Toronto, Canada

^e Division of Infectious Diseases, Department of Medicine, Sunnybrook Health Sciences Centre, University of Toronto, Canada

ARTICLE INFO

Keywords:

Cryptococcal meningitis
Pregnancy
Treatment
HIV
Malawi

ABSTRACT

This case reports cryptococcal meningitis in an HIV positive woman on antiretroviral therapy, presenting with left middle cerebral artery stroke at 30 weeks gestation. The patient had well-controlled HIV (CD4 count over 200 cells/mL). The immunosuppressive effects of the pregnancy likely contributed to the development of cryptococcal disease. The patient was successfully treated with two weeks of amphotericin B followed by fluconazole, delivered a healthy baby, but remained with a permanent severe neurological deficit.

1. Introduction

This case reports the difficulties in treating pregnant patients diagnosed with cryptococcal meningitis. Cryptococcal meningitis is a common disease in sub-Saharan Africa and is estimated to cause more than 100,000 deaths annually in the developing world [1,2], with mortality rates up to 68% [3]. It usually presents with a gradually progressive headache, and commonly results in cranial nerve defects and seizures. Worldwide cryptococcal meningitis is a disease occurring predominantly in those with T-cell immunosuppression secondary to HIV infection, and is an AIDS defining illness. We present a case here of a pregnant patient with an unusual presentation of cryptococcal meningitis.

2. Case

A 34-year-old female HIV-positive pregnant patient presented at 30 weeks gestation with collapse after feeling dizzy, with subsequent aphasia. There were no witnessed seizures, and she had not reported headache prior to collapse. The patient's HIV had been diagnosed 3 years prior, and she had been started at this time on antiretroviral therapy consisting of tenofovir, lamivudine and efavirenz, and co-trimoxazole prophylaxis, with excellent adherence evidenced by family report and medical records. There was no history of hypertension, diabetes, epilepsy, tuberculosis, or previous cryptococcal infection.

On examination, she had a fever of 38.2 °C, reduced movement of the right arm and leg, and a right-sided 7th cranial nerve palsy. Modified Rankin score 5/6. Glasgow coma score 10/15. Cardiovascular, respiratory, abdominal and fundoscopic examinations were unremarkable.

The patient was therefore diagnosed clinically with a left middle cerebral artery stroke. Differential diagnoses included infections such as toxoplasmosis, tuberculosis, cryptococcal meningitis, cerebral bacterial abscess, neurosyphilis, viral encephalitis, HIV encephalopathy, progressive multifocal leukoencephalopathy and cerebral malaria. Non-infective differentials included thromboembolic stroke, primary cerebral lymphoma, primary brain cancer or metastatic brain cancer.

Cerebrospinal fluid (CSF) from day 0 was positive for cryptococcus on India ink microscopy and cryptococcal antigen test (CrAg). Lumbar puncture was repeated on day +2 when the CrAg and India ink remained positive. The opening pressure on day +2 was 25 cm (normal range < 20 cm), white cell count 2 cells/μL (normal range 0–5 cells/μL), glucose 1.1 mmol/L (normal range 2.22–3.88), protein 1.09 g/L (normal range 0.15–0.40), and cryptococcal culture quantitative count was 45 colony forming units (cfu)/mL.

CD4 count on day +1 was 243 cells/mm³, HIV viral load on day +2 was undetectable, syphilis RPR was negative, hepatitis B surface antigen was negative, malaria rapid diagnostic test was negative, and creatinine/potassium levels were normal.

Echocardiogram on day +1 revealed no clots or other

* Corresponding author.

E-mail address: philip.bright@doctors.org.uk (P.D. Bright).

<http://dx.doi.org/10.1016/j.mmcr.2017.10.002>

Received 22 September 2017; Accepted 27 October 2017

Available online 28 October 2017

2211-7539/© 2017 Published by Elsevier B.V. on behalf of International Society for Human and Animal Mycology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

abnormalities. Chest X-ray was normal. Computerised tomography and magnetic resonance brain imaging were not available on site.

The patient received Ceftriaxone 2 g intravenously once daily to treat bacterial meningitis. This was stopped on day +1 following positive CSF tests for cryptococcal meningitis. She was started on aspirin 75 mg orally once/day. Anti-retroviral treatment and co-trimoxazole prophylaxis were continued. Two weeks of intravenous amphotericin B (AmB) 1 mg/kg once/day was started on day +2. Two litres of intravenous normal saline with 20 mmol potassium chloride, 1200 mg twice daily of oral potassium chloride, and three tablets once daily of magnesium glycerophosphate were given on each day of AmB treatment. Dexamethasone treatment (12 mg intravenously twice daily for 2 days) was given on days +3 and +4 to mature the foetal lungs, in view of the chance of premature delivery. The patient received a unit of blood on day +8 following a drop in haemoglobin from 9.1 g/dL on day +1 to 7.0 g/dL on day +6 (haemoglobin normal range 11.8–14.8 g/dL).

Lumbar puncture was repeated on day +7, when the opening pressure was 45 cm water, and the cryptococcal quantitative count was 30 cfu/mL. A final lumbar puncture was performed on day +14 when the opening pressure was 22, and the cryptococcal quantitative count was 0 cfu/mL.

The patient improved with the two weeks of AmB treatment, and oral fluconazole 400 mg twice per day then started. The fluconazole dose was reduced to 400 mg orally once/day 2 weeks later. The patient spontaneously entered labour 6 weeks following her presentation at 36 weeks gestation and delivered a healthy, but slightly underweight, baby boy (weight = 1.8 kg) without any noticeable congenital abnormalities.

An MRI Brain was performed after discharge (day +59) and is displayed in Fig. 1. Brain imaging was not performed earlier due to the clinical risk to the patient and her baby of problems arising during transport to another hospital 1 hour away. Brain imaging was still thought helpful to exclude differential diagnoses (e.g. brain abscess, tuberculoma, toxoplasmosis, lymphoma etc).

The patient was reviewed in clinic on day +123. She had gradually been improving at home, but severe neurological impairment persisted. At review her right-sided hemiparesis remained evident with contractures of the right foot and hand. She could walk with the assistance of one person, was able to eat and drink herself, and was breastfeeding. At this time she remained on fluconazole 200 mg once daily, tenofovir/lamivudine/efavirenz once daily, co-trimoxazole 960 mg once daily and aspirin 75 mg once daily.

3. Discussion

Experience on the treatment of patients with cryptococcal meningitis, disseminated cryptococcosis and pulmonary cryptococcosis in pregnancy is limited to case reports and small series totaling 50 cases

(Table 1). There have been only 3 documented cases of mother-to-child transmission of cryptococcus [4–6]. Three of the 8 (38%) cases reported of cryptococcal meningitis in confirmed HIV positive patients resulted in maternal death (Table 1).

Pregnancy may well be a predisposing factor for the development of cryptococcal disease [8]. Protection from cryptococcal disease is dependent on a good Th1 weighted cell-mediated immune response. The predisposition to cryptococcal disease in pregnancy likely relates to the necessity for tolerance by the mother's adaptive immune system to paternal antigens present in the foetus. CD4+ T-helper responses and number are reduced during pregnancy, with the greatest suppression in responses in the third trimester [9]. Additionally, in the context of patients with HIV infection off antiretroviral treatment, pregnancy reversibly reduces CD4 counts [10]. It is also possible to demonstrate a reduction in delayed hypersensitivity skin responses in pregnancy [7]. The weighting of T-cell responses in pregnancy changes from a Th1 cell-mediated (IFN- γ , IL-12) to a Th2 (IL-4, IL-10) antibody-mediated and T-regulatory (TGF- β) response [11]. The number of T-regulatory cells in peripheral blood actually increases during early pregnancy, peaks during the second trimester, and then declines post-partum [11]. Oestrogen reduces the activation of T-cells [12] and T-regulatory cell numbers correlate with oestrogen levels [13]. Progesterone is also considered to be immunosuppressive [14]. Progesterone has a positive role in the production of the T-regulatory cytokine IL-10, can induce Th2 weighted immune responses, reduce inflammatory cytokine production, reduce Th1 weighted immune responses and suppress allo-geneic responses [11]. Therefore pregnancy-related hormonal increases in both progesterone and oestrogen appear immunosuppressive.

Table 1 demonstrates that symptoms of cryptococcosis began pre-pregnancy in 2 cases (4%), during the first trimester in 8 cases (16%), the second trimester in 18 cases (36%), the third trimester in 18 cases (36%) and post-partum in 4 cases (8%). Symptomatic presentation therefore appears to be most common in the latter stages of pregnancy in keeping with the increased immunosuppression at this time.

This patient presented with a left middle cerebral artery stroke. Seventeen of 87 (20%) HIV positive cryptococcal meningitis patients had cortical or lacunar infarction [2], and 7 of 37 (19%) of immunocompetent patients with cryptococcal meningitis had cerebral infarction [15]. Therefore stroke is not an uncommon component of cryptococcal meningitis presentation.

Options for the treatment of cryptococcal disease include Amphotericin B based intravenous therapy, oral flucytosine and oral fluconazole. The Infectious Diseases Society of America (IDSA) guidelines on the treatment of disseminated cryptococcal disease in pregnancy recommends AmB based therapy with or without flucytosine [16]. AmB is a category B pregnancy drug indicating that there is no evidence of pregnancy risk in humans from its use following extensive clinical use for both cryptococcosis and other infections [17]. Flucytosine is a category C pregnancy drug indicating that not enough research




Fig. 1. MRI Brain Imaging of the patient at day +59. T2-weighted images (A, B) and DWI image (C) demonstrate extensive hyperintensity in the left frontal, parietal and temporal lobes, as well as the left caudate and lentiform nucleus, in keeping with a large left middle cerebral artery territory infarct. There is ventricular dilatation indicating hydrocephalus.

Download English Version:

<https://daneshyari.com/en/article/8484759>

Download Persian Version:

<https://daneshyari.com/article/8484759>

[Daneshyari.com](https://daneshyari.com)