

Accepted Manuscript

Title: Effect of different levels of pomegranate marc with or without polyethylene glycol on performance, nutrients digestibility and protozoal population in growing lambs

Authors: Ali Hatami, Daryoush Alipour, Fardin Hozhabri, Meysam Tabatabaei



PII: S0377-8401(17)31401-3
DOI: <https://doi.org/10.1016/j.anifeedsci.2017.11.004>
Reference: ANIFEE 13882

To appear in: *Animal Feed Science and Technology*

Received date: 3-5-2016
Revised date: 2-11-2017
Accepted date: 6-11-2017

Please cite this article as: Hatami, Ali, Alipour, Daryoush, Hozhabri, Fardin, Tabatabaei, Meysam, Effect of different levels of pomegranate marc with or without polyethylene glycol on performance, nutrients digestibility and protozoal population in growing lambs. *Animal Feed Science and Technology* <https://doi.org/10.1016/j.anifeedsci.2017.11.004>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Effect of different levels of pomegranate marc with or without polyethylene glycol on performance, nutrients digestibility and protozoal population in growing lambs

Ali Hatami¹, Daryoush Alipour^{1*}, Fardin Hozhabri², Meysam Tabatabaei^{3,4}

¹ Department of Animal Science, Faculty of Agriculture, Bu-Ali Sina University, Hamedan, Iran

² Department of Animal Science, Faculty of Agriculture, Razi University, Kermanshah, Iran

³ Biofuel Research Team (BRTeam), Karaj, Iran

⁴ Microbial Biotechnology and Biosafety Department, Agricultural Biotechnology Research Institute of Iran (ABRII), Karaj, Iran

* Corresponding author Tel: +98 8134425402

E-mail addresses: Alipourd@basu.ac.ir, Daryoush.alipour@gmail.com

Highlights

- Feeding pomegranate marc to growing lamb at low level (80 g/kg DM of diet) has no adverse effect
- Polyethylene glycol alleviate some negative nutritional effect of pomegranate marc at high level of feeding (160 g/kg DM of diet)
- Tannins in pomegranate marc shifted the nitrogen excretory pattern from urine to feces

Download English Version:

<https://daneshyari.com/en/article/8491059>

Download Persian Version:

<https://daneshyari.com/article/8491059>

[Daneshyari.com](https://daneshyari.com)