Grass carp fed a fishmeal-free extruded diet showed higher weight gain and nutrient utilization than those fed a pelleted diet at various feeding rates

Xiao-Qin Li, Huai-Bing Xu, Wen-Tong Sun, Xiao-Ying Xu, Zhen Xu, Xiang-Jun Leng

PII: S0044-8486(18)30163-7
DOI: doi:10.1016/j.aquaculture.2018.04.058
Reference: AQUA 633222

To appear in: *aquaculture*

Received date: 23 January 2018
Revised date: 28 April 2018
Accepted date: 30 April 2018

Please cite this article as: Xiao-Qin Li, Huai-Bing Xu, Wen-Tong Sun, Xiao-Ying Xu, Zhen Xu, Xiang-Jun Leng , Grass carp fed a fishmeal-free extruded diet showed higher weight gain and nutrient utilization than those fed a pelleted diet at various feeding rates. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Aqua(2017), doi:10.1016/j.aquaculture.2018.04.058

Grass carp fed a fishmeal-free extruded diet showed higher weight gain and nutrient utilization than those fed a pelleted diet at various feeding rates

Xiao-Qin Li[1,2,3], Huai-Bing Xu[1], Wen-Tong Sun[1], Xiao-Ying Xu[1], Zhen Xu[1], Xiang-Jun Leng[1,2,3*]

[1] National Demonstration Center for Experimental Fisheries Science Education (Shanghai Ocean University), Shanghai 201306, China

[2] Center for Research on Environmental Ecology and Fish Nutrition (CREEFN) of the Ministry of Agriculture, Shanghai Ocean University, Shanghai 201306, China

[3] Shanghai Collaborative Innovation Center for Aquatic Animal Genetics and Breeding, Shanghai 201306, China

***Corresponding author**: Xiang-Jun Leng, Tel: +86 21 61908760.

E-mail address: xjleng@shou.edu.cn (X.J. Leng).

Address: Hucheng Ring road 999, Lingang New City, Shanghai 201306, China.

Running title: Pelleted and extruded feeds, feeding rates and grass carp growth

[1] The first author is Xiaoqin Li, the co-first author is Huaibing Xu. These authors contribute equally to this work.