Tadpoles of Rhinella schneideri as reservoirs of trichodinids in continental aquaculture

G. Pala, G.M.R. Valladão, L.O. Alves, F. Pilarski, E.G. Lux Hoppe

PII: S0044-8486(17)31716-7
DOI: https://doi.org/10.1016/j.aquaculture.2018.01.017
Reference: AQUA 633019

To appear in: *aquaculture*

Received date: 24 August 2017
Revised date: 7 January 2018
Accepted date: 11 January 2018

Please cite this article as: G. Pala, G.M.R. Valladão, L.O. Alves, F. Pilarski, E.G. Lux Hoppe , Tadpoles of Rhinella schneideri as reservoirs of trichodinids in continental aquaculture. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Aqua(2017), https://doi.org/10.1016/j.aquaculture.2018.01.017

Tadpoles of *Rhinella schneideri* as reservoirs of trichodinids in continental aquaculture

G. Pala[1].; G.M.R. Valladão[2].; L.O. Alves[2]; F. Pilarski[2]; E.G. Lux Hoppe[1]

[1]Departamento de Medicina Veterinária Preventiva, Universidade Estadual Paulista (UNESP), Jaboticabal, São Paulo, Brazil.

[2]Aquaculture Center (CAUNESP), Universidade Estadual Paulista (UNESP), Jaboticabal, São Paulo, Brazil.

Email: hoppe@fcav.unesp.br