

Accepted Manuscript

Substitution of dietary fishmeal by soybean meal with inosine administration influences growth, digestibility, immunity, stress resistance and gut morphology of juvenile amberjack *Seriola dumerili*

Sakhawat Hossain, Shunsuke Koshio, Manabu Ishikawa, Saichiro Yokoyama, Nadia Mahjabin Sony, Jakiul Islam, Mayumi Maekawa, Takeshi Fujieda



PII: S0044-8486(17)30840-2
DOI: <https://doi.org/10.1016/j.aquaculture.2018.01.037>
Reference: AQUA 633039
To appear in: *aquaculture*
Received date: 1 May 2017
Revised date: 22 January 2018
Accepted date: 23 January 2018

Please cite this article as: Sakhawat Hossain, Shunsuke Koshio, Manabu Ishikawa, Saichiro Yokoyama, Nadia Mahjabin Sony, Jakiul Islam, Mayumi Maekawa, Takeshi Fujieda, Substitution of dietary fishmeal by soybean meal with inosine administration influences growth, digestibility, immunity, stress resistance and gut morphology of juvenile amberjack *Seriola dumerili*. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Aqua*(2017), <https://doi.org/10.1016/j.aquaculture.2018.01.037>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Substitution of dietary fishmeal by soybean meal with inosine administration influences growth, digestibility, immunity, stress resistance and gut morphology of juvenile amberjack *Seriola dumerili*

Md. Sakhawat Hossain ^{a,b,c*}, Shunsuke Koshio ^{a,b}, Manabu Ishikawa ^{a,b}, Saichiro Yokoyama ^a, Nadia Mahjabin Sony ^b, Md. Jakiul Islam ^d, Mayumi Maekawa ^e, Takeshi Fujieda ^e

^a*Laboratory of Aquatic Animal Nutrition, Faculty of Fisheries, Kagoshima University, Shimoarata 4-50-20, Kagoshima 890-0056, Japan*

^b*The United Graduate School of Agricultural Sciences, Kagoshima University, Korimoto 1-21-24, Kagoshima 890-0065, Japan*

^c*Department of Aquaculture, Faculty of Fisheries, Sylhet Agricultural University, Sylhet-3100, Bangladesh*

^d*Department of Fisheries Technology and Quality Control, Faculty of Fisheries, Sylhet Agricultural University, Sylhet-3100, Bangladesh*

^e*Research Institute for Bioscience Products & Fine Chemicals, Ajinomoto Co., Inc., Kawasaki, Kanagawa, 210-8681, Japan*

***Corresponding Author**

Md. Sakhawat Hossain, *Ph.D*

Laboratory of Aquatic Animal Nutrition

Faculty of Fisheries, Kagoshima University

Shimoarata 4-50-20; Kagoshima City; Kagoshima 890-0056, Japan

Phone: +81 99 286 4182; Hand phone: +81 080 3900 5285

Email: fishsakhawat@yahoo.com

Download English Version:

<https://daneshyari.com/en/article/8493397>

Download Persian Version:

<https://daneshyari.com/article/8493397>

[Daneshyari.com](https://daneshyari.com)