



Short communication

## Characterization of pantothenic acid deficiency and the dietary requirement of juvenile hybrid striped bass, *Morone chrysops* × *M. saxatilis*



Thiago Raggi, Alejandro Buentello, Delbert M. Gatlin III \*

Department of Wildlife and Fisheries Sciences, Texas A&M University System, College Station, TX 77843-2258, USA

### ARTICLE INFO

#### Article history:

Received 11 August 2015

Received in revised form 21 September 2015

Accepted 24 September 2015

Available online 26 September 2015

#### Keywords:

Hybrid striped bass  
Pantothenic acid  
Dietary requirement  
Deficiency

### ABSTRACT

An 8-week feeding trial was conducted to characterize the pantothenic acid deficiency signs and quantify the dietary requirement of juvenile hybrid striped bass, *Morone chrysops* × *Morone saxatilis*. A basal diet composed of fish muscle and crystalline amino acids was analyzed to contain 5 mg of pantothenate/kg and supplemented with calcium D-pantothenate resulting in five diets with graded pantothenate levels of 5, 10, 20, 30 and 40 mg/kg diet. Each diet was fed to triplicate groups of hybrid striped bass juveniles (initial average weight 1.6 g) in a recirculating system at  $26.0 \pm 1$  °C. Fish fed the unsupplemented basal diet performed poorly in terms of growth parameters and exhibited typical signs of pantothenic acid deficiency such as hemorrhages, sluggishness, high mortality, anemia and severe hyperplasia of the epithelial cells of gill lamellae. Fish fed the diet containing 10 mg of pantothenate/kg diet performed significantly better than fish fed the basal diet but poorly when compared with the responses of fish fed diets containing 20, 30 and 40 mg of calcium D-pantothenate/kg diet. Also, pantothenic acid-deficiency signs were apparent in fish fed the diet containing 10 mg/kg diet; whereas, no deficiency signs were observed in fish fed diets supplemented with the higher levels. Based on these data, the dietary pantothenic acid requirement of hybrid striped bass was quantified at 18.8 mg/kg diet based on broken-line regression analysis of weight gain data.

#### Statement of relevance:

Pantothenic acid deficiency in hybrid striped bass was characterized and the minimum dietary pantothenic acid requirement was determined to be 18.8 mg/kg based on weight gain. This information will assist in refining diet formulations for hybrid striped bass.

© 2015 Elsevier B.V. All rights reserved.

### 1. Introduction

Pantothenic acid (PA) is a water-soluble vitamin that functions as a part of the coenzyme A molecule in the metabolic release of energy from all three energy-providing nutrients, carbohydrate, fat and protein, by way of the tricarboxylic acid (TCA) cycle (NRC, 2011). Dietary essentiality of this nutrient has been established in numerous animals including various fish species (NRC, 2011). Response parameters used for PA requirement estimates are often based on the absence of deficiency signs, normal growth rate, liver tissue saturation and/or metabolic biomarker responses. The sign most frequently reported in fish caused by PA deficiency is severe hyperplasia of the epithelial cells of gill lamellae (clubbed gills), as well as hemorrhages, sluggishness, high mortality, and anemia (NRC, 2011). Due to this vitamin's

essentiality in the normal growth and health of fish, various researchers have attempted to determine qualitative and quantitative dietary requirements for PA which have ranged from 10 to 45 mg PA/kg diet (Cho and Woodward, 1990; Masumoto et al., 1994; Masumoto, 2002; Murai and Andrews, 1979; Ogino, 1967; Roem et al., 1991; Shimeno, 1991; Soliman and Wilson, 1992; Wen et al., 2009; Wilson et al., 1983).

Hybrid striped bass, crosses between white bass, *Morone chrysops*, and striped bass, *Morone saxatilis*, have shown considerable potential for aquaculture (Smith et al., 1985). The hybrids have desirable characteristics such as rapid growth and wide salinity tolerance (Smith et al., 1986). Some nutritional requirements have been determined for these hybrids (Brown et al., 1993; Keembiyehetty and Gatlin, 1992; NRC, 2011), but not that for PA, which may assist in the development of suitable diets. The present study was conducted to characterize PA deficiency signs and quantify the dietary requirement of juvenile hybrid striped bass, by evaluating growth performance and histopathological responses.

\* Corresponding author at: Department of Wildlife and Fisheries Sciences, Texas A&M University System, 2258 TAMU, College Station, TX 77843-2258, USA.  
E-mail address: [d-gatlin@tamu.edu](mailto:d-gatlin@tamu.edu) (D.M. Gatlin).

## 2. Materials and methods

### 2.1. Experimental diets

Semi-purified diets were formulated to provide 35% crude protein from solvent-extracted red drum (*Sciaenops ocellatus*) muscle and a crystalline amino acid premix. The red drum muscle was prepared in this laboratory by lyophilizing muscle tissue from adult red drum, as well as extracting three times in a boiling solution of hexane:ethanol (4:1 v/v) which reduced the endogenous lipid level so that it contributed approximately 0.65% lipid to the diet and 22.5% crude protein (Craig and Gatlin, 1997). The crystalline amino acid premix was formulated to provide the remaining 12.5% crude protein. Menhaden fish oil was supplemented to the lipid in the red drum muscle to provide a total of 9.7% dietary lipid.

The basal diet (Table 1) was analyzed to contain pantothenate at 5 mg/kg, and thus supplemental levels of calcium D-pantothenate (92% PA active) were adjusted to achieve total pantothenate levels of 10, 20, 30 and 40 mg/kg diet in the experimental diets. Samples of the experimental diets were sent to Eurofins Scientific (Des Moines, IA) for microbial analysis of PA according to AOAC (1990) procedures.

Preparation of the diets began with all dry components being individually weighed into 3.8-L plastic bags. Once all dry ingredients had been weighed, diets were thoroughly mixed using a commercial V-mixer for 30 min. Diets were then placed in a commercial Hobart mixer where the fish oil and water were added. Diets were then pressure pelleted through a 3-mm die using a Hobart meat grinder. Pellets were then crumbled by hand and dried at 25 °C. After drying, pellets were screened to remove fines and stored at –20 °C until fed.

### 2.2. Experimental procedures

Juvenile hybrid striped bass were obtained from Keo Fish Farms (Lonoke, Arkansas) and transported by truck to the Texas A&M Aquacultural Research and Teaching Facility. Prior to the beginning of the feeding trial, fish underwent a 1-week conditioning period to adjust to a purified basal diet and standardized experimental conditions. Each of the five dietary treatments was randomly assigned to triplicate 110-L aquaria in a recirculating system at the Texas A&M University Aquacultural Research and Teaching Facility. Each aquarium contained 15 fish with an initial average weight of 1.6 g per fish. Water flow through each culture chamber was maintained at approximately 0.65 L/min via a 1.5 HP pump that recirculated water through biological and mechanical filters to maintain appropriate water quality (total ammonia nitrogen < 0.6 mg/L). Salinity was maintained at 6.5 to 8‰ using well water and synthetic sea salt (Fritz Industries Inc., Dallas, TX, USA). Low-pressure electrical blowers provided aeration via air stones and constantly maintained dissolved

oxygen (DO) levels at or near saturation. Water temperature was regulated by controlling ambient air temperature and remained at  $26 \pm 1$  °C throughout the trial. A daily light:dark cycle of 12:12 h was provided by fluorescent lighting.

Fish were fed each diet at 8% of body weight per day for the first week, and then feeding rate was progressively reduced to minimize overfeeding while maintaining a level close to satiation. Fish were weighed every week and the amount of diet fed was adjusted accordingly. Dead fish were removed and not replaced during the experiment. Fish were fed the experimental diets for 8 weeks.

### 2.3. Growth performance and histopathology

Percentage of body weight gain (WG) [ $100 \times (\text{final body weight} - \text{initial body weight}) / \text{initial body weight}$ ], feed efficiency (FE) [ $(\text{final body weight} - \text{initial body weight}) / \text{feed intake}$ ], and survival [ $100 \times (\text{final fish number} / \text{initial fish number})$ ] were calculated.

Fifteen fish were randomly selected from the initial group before assignment to the various diets and fixed in 10% neutral buffered formalin to serve as control histological samples. At the end of the feeding trial, gills from fish fed the diet supplemented with 15 mg PA/kg and the basal diet were dissected, placed in 10% neutral buffered formalin, embedded in Paraplast (Lancer Co., St. Louis, MO), sectioned and stained with hematoxylin and eosin. The gill sections were examined microscopically for histological changes.

### 2.4. Statistical analysis

All data were subjected to analysis of variance and Duncan's multiple-range test (Duncan, 1955) to determine differences in treatment means ( $P < 0.05$ ). Weight gain data also were analyzed by least-squares regression using the broken-line model (Robbins et al., 2006) to estimate the minimum dietary PA requirement of hybrid striped bass for optimal growth. Data analyses were performed using the SAS/PC statistical software (SAS Inst. Inc., Cary, NC).

## 3. Results and discussion

The importance of dietary PA for hybrid striped bass was evident in the present study. The response of fish to increasing levels of dietary calcium D-pantothenate is presented in Table 2. Significantly reduced weight gain, anemia, anorexia, low survival, and sluggishness were observed in fish fed the basal diet. Feed efficiency and percent survival increased stepwise up to a dietary level of 15 mg calcium D-pantothenate/kg diet and plateaued. Dietary pantothenic acid had no effect on condition factor of the fish.

Marked interlamellar proliferative lesions were observed in the gills of fish fed the basal diet. The lesions were visible as early as week 3 and were most pronounced at the distal end of the gill filaments giving a

**Table 1**  
Composition of the basal diet.

Ingredient	% (dry weight)
Red drum muscle (lipid extracted)	21.9
Amino acid mix <sup>a</sup>	12.5
Dextrin <sup>b</sup>	30.0
Menhaden oil <sup>c</sup>	9.7
Vitamin premix <sup>d</sup>	3.0
Mineral premix <sup>d</sup>	4.0
Carboxymethylcellulose <sup>b</sup>	2.0
Celufil <sup>b</sup>	14.4
CaHPO <sub>4</sub> ·2H <sub>2</sub> O <sub>2</sub>	1.0
Aspartate/glutamate premix <sup>b</sup>	1.5
Calcium D-pantothenate <sup>b</sup>	0.0

<sup>a</sup> Contains (g/100 g): arginine 9.44; glycine 11.92; histidine 3.76; isoleucine 6.16; leucine 9.60; methionine 4.08; phenylalanine 2.16; tyrosine 4.32; serine 2.16; threonine 7.04; tryptophan 1.92; valine 5.84; proline 15.76; and alanine 15.84.

<sup>b</sup> US Biochemical Corp., Cleveland, OH, USA.

<sup>c</sup> Omega Protein Corporation, Inc. Houston, TX, USA.

<sup>d</sup> Same as Moon and Gatlin (1991).

**Table 2**  
Responses of fingerling hybrid striped bass to graded levels of dietary calcium D-pantothenate levels<sup>1,2</sup>.

Dietary pantothenate mg/kg diet	Initial weight g/group	Weight gain % of initial wt.	Feed efficiency g gain/g feed	Survival %
5	19.3 ± 1.4	212 ± 40.7 <sup>c</sup>	0.35 ± 0.04 <sup>c</sup>	17.8 ± 8.0 <sup>b</sup>
10	20.2 ± 1.5	628 ± 35.0 <sup>b</sup>	0.54 ± 0.05 <sup>b</sup>	75.6 ± 24.4 <sup>a</sup>
20	20.6 ± 0.6	1358 ± 57.2 <sup>a</sup>	0.75 ± 0.01 <sup>a</sup>	97.8 ± 2.2 <sup>a</sup>
30	20.1 ± 1.3	1373 ± 25.1 <sup>a</sup>	0.75 ± 0.01 <sup>a</sup>	100 ± 0.0 <sup>a</sup>
40	20.8 ± 1.6	1338 ± 93.3 <sup>a</sup>	0.75 ± 0.03 <sup>a</sup>	100 ± 0.0 <sup>a</sup>

<sup>1</sup> Values are means ± standard error of three groups of fish fed the same experimental diet ( $n = 3$ ) with 15 fish per group.

<sup>2</sup> Numbers with different superscripts in the same column indicate significant ( $P \leq 0.05$ ) differences according to Duncan's multiple-range test.

Download English Version:

<https://daneshyari.com/en/article/8494171>

Download Persian Version:

<https://daneshyari.com/article/8494171>

[Daneshyari.com](https://daneshyari.com)