

Accepted Manuscript

Low-value fish used as feed in aquaculture were a source of furunculosis caused by atypical *Aeromonas salmonicida*

S.-Y. Choi, C.-S. Kim, M.-J. Oh, H.-D. Jeong, D.-H. Kim

PII: S0044-8486(13)00235-4
DOI: doi: [10.1016/j.aquaculture.2013.05.014](https://doi.org/10.1016/j.aquaculture.2013.05.014)
Reference: AQUA 630661

To appear in: *Aquaculture*

Received date: 18 April 2013
Revised date: 10 May 2013
Accepted date: 14 May 2013



Please cite this article as: Choi, S.-Y., Kim, C.-S., Oh, M.-J., Jeong, H.-D., Kim, D.-H., Low-value fish used as feed in aquaculture were a source of furunculosis caused by atypical *Aeromonas salmonicida*, *Aquaculture* (2013), doi: [10.1016/j.aquaculture.2013.05.014](https://doi.org/10.1016/j.aquaculture.2013.05.014)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Low-value fish used as feed in aquaculture were a source
of furunculosis caused by atypical *Aeromonas
salmonicida***

S.-Y. Choi^a, C.-S. Kim^a, M.-J. Oh^a, H.-D. Jeong^b, D.-H. Kim^{a,*}

^aDepartment of Aqualife Medicine and Fish Health Center, Chonnam National University, Yeosu 550-749, Republic of Korea

^bDepartment of Aquatic Life Medicine, Pukyong National University, Busan 608-737, Republic of Korea

*Corresponding author. Tel.: +82 61 659 7177.

E-mail address: kimdh@chonnam.ac.kr (D.-H. Kim).

Download English Version:

<https://daneshyari.com/en/article/8495545>

Download Persian Version:

<https://daneshyari.com/article/8495545>

[Daneshyari.com](https://daneshyari.com)