

## Accepted Manuscript

Effects of dietary rapeseed meal on laying performance, egg quality, apparent metabolic energy, and nutrient digestibility in laying hens

L.P. Zhu , J.P. Wang , X.M. Ding , S.P. Bai , Q.F. Zeng ,  
Z.W. Su , Y. Xuan , K.Y. Zhang

PII: S1871-1413(18)30183-5  
DOI: [10.1016/j.livsci.2018.06.007](https://doi.org/10.1016/j.livsci.2018.06.007)  
Reference: LIVSCI 3480



To appear in: *Livestock Science*

Received date: 18 October 2017  
Revised date: 13 June 2018  
Accepted date: 15 June 2018

Please cite this article as: L.P. Zhu , J.P. Wang , X.M. Ding , S.P. Bai , Q.F. Zeng , Z.W. Su , Y. Xuan , K.Y. Zhang , Effects of dietary rapeseed meal on laying performance, egg quality, apparent metabolic energy, and nutrient digestibility in laying hens, *Livestock Science* (2018), doi: [10.1016/j.livsci.2018.06.007](https://doi.org/10.1016/j.livsci.2018.06.007)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

### Highlights

- Low glucosinolate rapeseed meal content in hens' diet should be lower than 117.6 g/kg.
- Egg weight decreased when rapeseed meal intake was more than 13.6 g/hen/day.
- Egg production decreased when rapeseed meal intake was more than 6.5 g/hen/day.

Download English Version:

<https://daneshyari.com/en/article/8501898>

Download Persian Version:

<https://daneshyari.com/article/8501898>

[Daneshyari.com](https://daneshyari.com)