


Comparison of two herbal essential oils, probiotic, and mannan-oligosaccharides on egg production, hatchability, serum metabolites, intestinal morphology, and microbiota activity of quail breeders

M. Hajiaghapour, V. Rezaeipour



PII: S1871-1413(18)30053-2
DOI: <https://doi.org/10.1016/j.livsci.2018.02.007>
Reference: LIVSCI3401

To appear in: *Livestock Science*

Received date: 5 August 2017
Revised date: 11 January 2018
Accepted date: 8 February 2018

Cite this article as: M. Hajiaghapour and V. Rezaeipour, Comparison of two herbal essential oils, probiotic, and mannan-oligosaccharides on egg production, hatchability, serum metabolites, intestinal morphology, and microbiota activity of quail breeders, *Livestock Science*, <https://doi.org/10.1016/j.livsci.2018.02.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Comparison of two herbal essential oils, probiotic, and mannan-oligosaccharides on egg production, hatchability, serum metabolites, intestinal morphology, and microbiota activity of quail breeders

M. Hajiaghapour, V. Rezaeipour*

Department of Animal Science, Qaemshahr Branch, Islamic Azad University, Qaemshahr, Iran

* Corresponding author.

E-mail address: vrezaeipour@qaemiau.ac.ir (V. Rezaeipour).

Download English Version:

<https://daneshyari.com/en/article/8502003>

Download Persian Version:

<https://daneshyari.com/article/8502003>

[Daneshyari.com](https://daneshyari.com)