

Accepted Manuscript

Use of urinary γ -glutamyl transferase (GGT) to monitor the pattern of proteinuria in dogs with leishmaniasis treated with N-methylglucamine antimoniate

Saverio Paltrinieri, Giulia Mangiagalli, Fabrizio Ibba



PII: S0034-5288(17)30896-2
DOI: doi:[10.1016/j.rvsc.2018.05.014](https://doi.org/10.1016/j.rvsc.2018.05.014)
Reference: YRVSC 3578
To appear in: *Research in Veterinary Science*
Received date: 24 August 2017
Revised date: 14 January 2018
Accepted date: 23 May 2018

Please cite this article as: Saverio Paltrinieri, Giulia Mangiagalli, Fabrizio Ibba , Use of urinary γ -glutamyl transferase (GGT) to monitor the pattern of proteinuria in dogs with leishmaniasis treated with N-methylglucamine antimoniate. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Yrvsc(2017), doi:[10.1016/j.rvsc.2018.05.014](https://doi.org/10.1016/j.rvsc.2018.05.014)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Use of urinary γ -glutamyl transferase (GGT) to monitor the pattern of proteinuria in dogs with leishmaniasis treated with N-methylglucamine antimoniate

Saverio Paltrinieri, DVM, PhD, Dipl. ECVCP^{a,b,*} saverio.paltrinieri@unimi.it, Giulia Mangiagalli, DVM^a, Fabrizio Ibba, DVM, PhD^{a,c}

^aDepartment of Veterinary Medicine, University of Milan, Milan, Italy

^bVeterinary teaching hospital, University of Milan

^cVeterinary Clinic Poggio dei Pini, Capoterra, Cagliari, Italy

*Corresponding author.

Abstract

The aim of this study was to assess if the coupled analysis of the urinary protein to creatinine (UPC) ratio and of the GGT/UC ratio (the ratio between urinary γ -glutamyl transferase activity and urinary creatinine) may be used in treated leishmaniotic dogs to differentiate dogs with transient impairment of tubular function from dogs with persistent tubular damage.

Download English Version:

<https://daneshyari.com/en/article/8503877>

Download Persian Version:

<https://daneshyari.com/article/8503877>

[Daneshyari.com](https://daneshyari.com)